

531 Rec'd PCT/P 22 JUN 2001

1/PRTS

HEAT TOLERANT BROCCOLI

Ins 01

FIELD OF THE INVENTION

- 5 This invention is in the field of plant breeding. In particular, this invention relates to the development of heat tolerant broccoli (*Brassica oleracea* L. var. *italica*).

BACKGROUND OF THE INVENTION

- 10 Broccoli (*Brassica oleracea* L. var. *italica*) has become an increasingly popular crop worldwide especially in health-conscious areas of the western world such as the North America, Europe, and Japan. An average broccoli stalk contains only 30 calories and provides 240% of the recommended daily allowance of vitamin C  
15 plus 10% of the recommended daily allowance of vitamin A. In addition to its nutritional value, some recent studies have shown that broccoli aids in the prevention of some forms of cancer.

- Broccoli is a cool weather crop. High temperatures ( $> 80^{\circ}\text{F}$ ) for even relatively short periods of time and warm temperatures  
20 ( $> 75^{\circ}\text{F}$ ) for extended periods of time cause broccoli heads to be rough with uneven flower bud sizes and thus commercially unacceptable. {(Björkman, T., et al. (1998) High temperature arrest of inflorescence development in broccoli (*Brassica oleracea* var. *italica* L.) Journal of Experimental Botany 49:101-106.} As a result  
25 of the high sensitivity to heat during growth, broccoli can only be grown in limited areas under cool weather conditions.

- Previous attempts at identifying heat tolerant broccoli cultivars have not been successful because broccoli is sensitive to relatively short periods of heat stress thereby making field observations too  
30 variable for effective genetic screening. Björkman, et al. (1998).

[illegible]

## SUMMARY OF THE INVENTION

10

15

invention are capable of germinating into a plant that is predominately mildew resistant.

20

25

30

to those seeds designated M7028, M7007, M7009, M7022, 393-2-19, H7008, H7022, 393-2-47, 98-2192, 98-2088, 98-2061, H7007, H7028, H7010, and H7021R. The broccoli seed of this

5 invention further include lines 4243, 4221, 4441, 4274-1, 4274-2,  
4278-1, 4284-1, 4285-1, 4354-1, 4354-2, 4377-1, 4318-1, 4320-  
1, 4320-2, 4321-1, 4437-1, 4476-1, 4462-1, 4308-2, 4309-1,  
4355-1, 4412-1, 4301, 4303, 4304, 4317, 4468, 4470, 4471,  
4263-1, 4430-1, 4450-1, 4450-2, 4432-1, 4267-1, 7861, 7864,  
10 7865, 7881, 7887, 7935, 8092, 7883, 7914, 7770, 7778.

The broccoli seed of this invention further include lines 4201, 4219, 4237, 4280, 4287, 4288, 4289, 4290, 4291, 4458-1, 4460-1, 4415, 4418, 4395-2.

Each of the lines of this invention can be crossed with other  
15 broccoli lines.

The broccoli seeds of this invention include inbred lines, hybrid lines, male lines and female lines, all of which are heat tolerant and capable of producing a commercially acceptable head under heat stress growth conditions.

20 This invention is further directed to broccoli plants or parts of  
broccoli plants produced from the broccoli seed of the invention.  
The invention is further directed to broccoli plants regenerated from  
tissue culture of the broccoli plants of this invention. The tissue  
culture of the invention comprises regenerable cells including  
25 meristematic tissue, anthers, leaves, ovules, roots, embryos,  
protoplasts and pollen and plants regenerated from these cells.

The invention is further directed toward transgenic heat tolerant broccoli plants. The transgenic heat tolerant broccoli lines may be resistant to various herbicides or pesticides.

30           The invention is further directed to broccoli plants having all of  
the phenotypic characteristics of the plants produced from the heat

tolerant broccoli seed of the invention. The invention is further directed to plants resulting from selecting, crossing, breeding or otherwise altering the broccoli plants of this invention.

5 The invention is further directed to biological material isolated from the plants of this invention. Such material includes but is not limited to RNA, DNA, protein and carbohydrate. The DNA of these plants includes the gene(s) involved in heat tolerance.

10 This invention is further directed to the seeds and plants produced from crossing other broccoli lines with plants grown from the seed of this invention.

This invention is further directed to methods of breeding heat tolerant broccoli lines.

#### **BRIEF DESCRIPTION OF THE DRAWINGS**

15 The invention will be better understood by reference to the drawings in which:

Figure 1 shows various broccoli head shapes. Broccoli heads are referred to as domed, semi-domed and deep-domed. The shapes of the various domes are (1) circular; (2) transverse broad elliptic; (3) 20 transverse elliptic and (4) transverse elliptic narrow.

Figure 2 shows a cross section of eight 40-inch beds utilized for field production of the hybrid seed of this invention. In this diagram: F = female line seed-line; M1 = first male planting seed-line; M2 = second male planting seed-line; M3 = third male planting seed- 25 line and M4 = fourth male planting seed-line.

#### **DETAILED DESCRIPTION OF THE INVENTION**

In order to more completely understand the invention, the following definitions are provided.

**Broccoli:** Broccoli (*Brassica oleracea* L. var. *italica*) is a cool season vegetable in the mustard family. Principal broccoli varieties currently grown in California include, in the coastal valleys, Everest, Greenbelt, Legacy, Marathon, Ninja, Olympia, Pinnacle, Pirate, Republic, Shogun, and Sultan; in the desert valleys, Arcadia, Captain, Emperor, Everest, Galaxy, Galleon, Greenbelt, Major, Marathon, Ninja, Packman, Patriot, Pirate, and Sultan; and in the San Joaquin Valley, Arcadia, Captain, Everest, Greenbelt, Legacy, Legend, Marathon, Pirate, and Republic. Varieties grown in the Pacific Northwest are: Arcadia, Emerald City, Excelsior, Pakman Patriot, Pirate, Regal, Arcadia, Buccaneer, Emerald City, Emperor, Everest, Excelsior, Green Belt, Green Valiant, Laguna, Legend, Liberty, Major, Marathon, Pakman, Patriot, Pinnacle, Pirate, Premium Crop, Regal, Shogun, Samurai, Triathlon, Windsor, Barbados, Embassy, Green Comet, Green Defender, HMX 1134, Idol. Because of heat sensitivity, broccoli is typically grown for harvest in the spring and fall.

**Commercially Acceptable Broccoli:** Commercially acceptable broccoli is broccoli which vegetable growers/shippers find acceptable for sale and consumers find acceptable for personal purchase and, ultimately, human consumption. Commercially acceptable broccoli has small uniform beads, good blue-green to green color, and tight, dome-shaped heads that extend above the leaves for ease of harvest. In commercial plantings under optimum conditions, large leafy broccoli plants produce a compact flower head on a tall, green, branching stalk. The center flower head is from 3 to 8 inches (7.5-20 cm) in diameter and plants average 24 to 36 inches (60-90 cm) tall. Hollow stems, water head rot, brown or yellow beads, bracts (leaflets) within heads, uneven bead size, and excessive branching

are undesirable and commercially unacceptable defects in broccoli that can be caused by exposure to heat.

**Heat Tolerant Broccoli:** Heat tolerant broccoli is broccoli that  
5 will produce a commercially acceptable product when grown under heat stress growth conditions for broccoli.

**Heat Stress Growth Conditions:** Heat stress growth conditions for broccoli are elevated temperature growth conditions that result in  
10 broccoli that exhibit heat stress symptoms that result in a commercially unacceptable product. Heat stress symptoms include non-uniform beads; brown, yellow, light-green or purple colored heads; loose flat heads; prominent leaflets that come through the broccoli head as bracts; hollow stems; water head rot and excessive  
15 branching.

**Single Plant Selection:** Single plant selection is the process of selecting single plants, which exhibit desired traits or characteristics. Seeds from the single plant are collected, stored and then grown in a  
20 subsequent growing period for further selections.

**Massed:** Broccoli plants are massed when a number of plants are selected and brought together for cross-pollination as a group. Massing prevents further inbreeding and tends to "fix" the broccoli  
25 line at the stage from which the selections were made.

**Self-Pollination/Self-Pollinator:** Self-pollination is the process of putting pollen from a plant onto a receptive female flower-part of that same plant. A plant that is a self-pollinator is a plant that  
30 accepts its own pollen to make seed that typically will give rise to

20250120063500

plants very similar or the same as the self pollinator plant. A plant that is self-pollinated is said to be selfed.

**Self-Incompatible:** A self-incompatible plant will not, under normal conditions, accept its own pollen nor generate any self-seed. Self-incompatible lines are generally designated "female." Self-incompatible lines are generally crossed with other lines to produce hybrid seed.

**Self-Compatible:** A self-compatible plant accepts its own pollen and will produce self-seed. Self-compatible lines are generally designated "male."

**Progeny:** Progeny is a broccoli line that is the offspring of the previous generation broccoli line.

**Sessile:** Attached to the stem by the base of the leaf.

**Petiolate:** Attached to the stem via a petiole.

**Hybrid:** The progeny of cross-fertilization between parents belonging to different genotypes.

**Hybrid Vigor:** The phenomenon in which the cross of two parent stocks produces hybrids that show increased vigor/heterosis compared to the parent stocks.

**Inbred Lines:** A nearly homozygous line produced by continuous inbreeding.

**Pedigree Bre ding:** A system of breeding in which individual plants are selected in the segregating generations from a cross on the basis of their desirability and on the basis of a pedigree record.

5           The terminology used to describe the broccoli plants of this invention are generally those used by the Plant Variety Protection Office in PVP form STD-470-44 "Objective Description of Variety Broccoli (*Brassica oleracea* var. *italica*).". The following terminology is used herein in comparative study #1 and comparative study #2.

10

**1. REGION OF ADAPTATION (Area where best adapted in USA):**

(1) Northwest; (2) NorthCentral; (3) Northeast; (4) Southeast; (5) Southwest; (6) Most regions and (7) Pacific Coast.

15

**2. MATURITY (Main Crop at 50% Harvest):**

20   Harvest Season: (1) Fall; (2) Fall/Winter; (3) Winter/Spring; (4) Spring/Summer; (5) Summer; and (6) Summer/Fall.

Spring Planted: (1) Days from Direct Seeding to 50% Harvest; (2) Days from Transplanting to 50% Harvest; and (3) Length of Harvest Period in days.

25

Fall Planted: (1) Days from Direct Seeding to 50% Harvest; (2) Days from Transplanting to 50% Harvest; and (3) Length of Harvest Period in days.

30   Time of beginning of flowering (50% of plants with at least 10% flowers: (1) Early; (2) Med-Early; (3) Medium; (4) Med-Late; and (5) Late.

35   **3. SEEDLING:**

Cotyledon Color: (1) Yellow-Green; (2) Light Green; (3) Medium Green; (4) Dark Green; (5) Blue-Green; and (6) Purple-Green.

40   Cotyledon Anthocyanin: (1) Absent; (2) Weak; (3) Intermediate; and (4) Strong.



Hypocotyl Anthocyanin: (1) Absent; (2) Weak; (3) Intermediate; and (4) Strong.

5

**4. PLANT (At Harvest):**

Plant Height: (cm) from soil line to top of leaves

10 Head Height: (cm) from soil line to top of head

Plant Branches: (1) Few; (2) Medium; and (3) Many.

Plant Habit: (1) Spreading; (2) Intermediate; and (3) Compact.

15

Market Class: (1) Fresh Market; (2) Processing; and (3) Both

Life Cycle: (1) Annual; (2) Biennial; and (3) Perennial.

20 Type of Variety: (1) Inbred; (2) Open-Pollinated; and (3) First generation Hybrid.

**5. OUTER LEAVES (at Harvest):**

25

Number of Leaves per Plant:

Width at midpoint of plant including petiole:

30 Length at midpoint of plant including petiole:

Petiole Length:

35 Leaf Ratio-Length/Width: (1) (2:1); (2) (3:1); (3) (4:1); (4) (5:1); and (5) (6:1).

Leaf Attachment: (1) Sessile; (2) Petiolate; and (3) Sessile and Petiolate (both).

40 Wax Presence: (1) None; (2) Weak; (3) Intermediate; and (4) Strong.

Foliage Color (with wax if present): 1 Light Green; (2) Medium Green; (3) Dark Green; (4) Grey-Green; (5) Blue-Green; and (6) Purple-Green.

45

Leaf Shape: (1) Narrow Elliptic; (2) Elliptic; and (3) Broad Elliptic.

Leaf Base: (1) Blunt and (2) Pointed.

Leaf Apex: (1) Blunt and (2) Pointed.

5

Leaf Margins: (1) Straight; (2) Slightly Wavy; and (3) Very Wavy.

Leaf Veins: (1) Thin; (2) Intermediate; and (3) Thick.

10 Midrib: (1) Not Raised; (2) Slightly Raised; and (3) Raised.

Blistering (1) None; (2) Weak; and (3) Intermediate; and (4) Strong.

15 Attitude (Leaf Angle from Ground): (1) Horizontal (0-15 degrees);  
(3) Semi-erect (35-55 degrees); and (5) Erect (80-100 degrees).

Torsion of Leaf Tip: (1) None; (2) Weak; (3) Intermediate; and (4) Strong.

20 Profile of Upper Side of Leaf: (1) Concave; (2) Planar; and (3) Convex.

#### 6. HEAD (At Market Maturity):

25

Diameter at widest point:

Depth:

30 Weight: market trimmed

Color: (1) Light Green; (2) Medium Green; (3) Dark Green; (4) Blue/Green; and (5) Purple.

35 Head Shape: (1) Circular; (2) Transverse Broad Elliptic; (3) Transverse Elliptic; and (4) Transverse Elliptic Narrow.

Dome Shape: (1) Domed; (2) Semi-domed; and (3) Deep Domed.

40 Head Size: (1) Small; (2) Medium; and (3) Large.

Compactness: (1) Long Pedicels (Loose); (2) Medium; and (3) Short Pedicels (Tight).

45 Surface Knobbling: (1) Fine; (2) Medium; and (3) Coarse.

Beads size: (1) Small; (2) Medium; and (3) Large.

Flower Buds: (1) Even in size; and (2) Uneven in size (cateye).

- 5 Anthocyanin Coloration: (1) Absent; 2 Present; (3) Leaf Axils; (4) Leaf Veins; (5) Leaf Blade; (6) Entire Plant; and (7) Leaf Petiole.

Color of Head Leaves: (1) White; (2) Green; (3) Red; and (4) Purple.

- 10 Secondary Heads: (1) Completely absent; (2) Basal; (3) Combination; and (4) Axillary along entire main stem up to main head.

Prominence of Secondary Heads: (1) Weak, (2) Intermediate; and (3) = Strong.

15

Number of Secondary Heads:

7. COLOR:

20

Flower Color: (1) White; (2) Cream; and (3) Yellow.

Flower Stalk Color: (1) Green; (2) Purple; and (3) Variegated.

25

8. DISEASE RESISTANCES:

1 = Most Susceptible

5 = Intermediate

9 = Most Resistant

30

Black Leg (*Leptosphaeria maculans*)

Black Spot (*Alternaria* spp.)

Black Rot

Bottom Rot (*Rhizoctonia solani*)

Cauliflower Mosaic Virus

Cerospora Leaf Spot (*Cercospora brassicicola*)

Clubroot (*Plasmodiophora brassicae*)

Downy Mildew (*Peronospora parasitica*)

Erwinia Sp.

Phytophthora Root Rot (*Phytophthora megasperma*)

Powdery Mildew (*Erysiphe polygoni*)

Pseudomonas

Ring Spot (*Mycosphaerella brassicicola*)

Black Leg

Black Spot

Black Rot

Bottom Rot

Cauliflower Mosaic Virus

Cerospora Leaf Spot

Clubroot

Downy Mildew

Erwinia Sp.

Phytophthora Root Rot

Powdery Mildew

Pseudomonas

Ring Spot

Turnip Yellow Mosaic Virus  
Verticillium wilt (*Verticillium albo-atrum*)  
White Blister (*Albugo candida*)  
Xanthomonas campestris  
Yellows (*Fusarium oxysporum*)

Turnip Yellow Mosaic Virus  
Verticillium wilt  
White Blister  
Xanthomonas campestris  
Yellows

**9. OTHER RESISTANCE:**

- 1 = Most susceptible  
5 = Intermediate  
5 9 = Most Resistant

Insect  
Buttoning  
Blindness  
Bolting  
Brown beads  
Drought  
Cold  
Hollow Stem  
Riceyness  
Whiptail

**10. HEAT TOLERANCE:**

- Heat tolerance was measured on a scale of 1-9 with 9 being the most heat tolerant and 1 being the least heat tolerant. For heat tolerance, ratings of five (5) or below are indicative of a broccoli plant that produced a commercially unacceptable head. A rating of six (6) is indicative of broccoli plants that exhibit no heat stress symptoms when exposed to 90°F. A rating of seven (7) is indicative of broccoli plants that exhibit no heat stress symptoms when exposed to 95°F. A rating of eight (8) is indicative of broccoli plants that exhibit no heat stress symptoms when exposed to 100°F. A rating of nine (9) is indicative of broccoli plants that exhibit no heat stress symptoms when exposed to 105°F. In some studies, plants were not exposed to a temperature of 105°F so that a rating of 9.0 could not be assigned.

Under some circumstances, the heat tolerant ratings are followed by a (+) or (-) to indicate a plant exhibiting symptoms

202502005850

slightly better (+) or slightly worse (-) than the assigned number. In other circumstances, the ratings are presented as a fraction of a rating number. For example, a rating of 7.1 is slightly better than a rating of 7.0. A rating of 7.5 is half way between a rating of 7.0 and 8.0. A rating of 6.8 is slightly worse than a rating of 7.0. A slightly higher rating means that the heat stress symptoms were slightly less evident. Similarly, a slightly lower rating means that the heat stress symptoms were less evident.

Taking into account these definitions, the present invention is directed to heat tolerant broccoli plants. The heat tolerant broccoli of this invention is capable of producing a commercially acceptable product when grown under heat stress conditions.

Heat stress is exhibited in broccoli by a number of different symptoms. These symptoms include non-uniform heads; brown, yellow, light-green or purple colored heads; flat heads; bracts (leaflets in the head); rapid fracturing of the head, which reduces the harvest period; "cateye" (death of growing points), extremely small heads, and hollow stems.

Each of these symptoms is generally viewed as commercially unacceptable. The greater the number of heat stress symptoms, the more commercially unacceptable the broccoli plant. Heat stress symptoms in broccoli result from a number of interacting factors. The most important of these interacting factors are the temperature, the duration of the high temperature exposure (hours, days, weeks), the available soil moisture supply and the wind speed. Of critical importance is the timing of the exposure to the heat stress conditions during the growth cycle of broccoli. It has been shown that heat stress of broccoli can be due to an inhibition of the enlargement of broccoli bud primordia. Broccoli buds are not as sensitive to heat once they differentiate. The different heat sensitivity and resulting contrast between the delayed buds and the

unaffected buds causes the uneven head appearance under heat stress growth conditions. If heat stress occurs prior to bud development (i.e., during vegetative development) no injury is generally seen. If the heat stress is applied late in bud development, many buds are affected but these buds can be obscured by the older buds.

Because of extreme sensitivity to heat stress, broccoli grown in the spring and the fall, when cooler temperatures are the norm, are at less risk of heat stress. However, a single day during the spring or fall with a high temperature of 100°F or several (2-3) warmer days (>80°F) or multiple (5-7) warm days (>75°F) at the critical point during broccoli bud development can render an entire field so damaged by heat stress that none of the heads are commercially acceptable.

The present invention is thus directed toward the development of heat tolerant broccoli varieties and hybrids. The broccoli varieties and hybrids of this invention will produce commercially acceptable heads when the plants are grown during heat stress growth conditions during late spring, summer, and early fall in California, Arizona, Mexico, and many other areas traditionally considered to be too warm for broccoli growth, or at risk of heat stress.

### **Broccoli Production**

Broccoli may be grown by transplant production or by direct seeding. For transplant production, plants may be started in hotbeds or greenhouses. Broccoli seedlings grown in a hotbed need a loose, easily pulverized loam that is not too fertile. If the plants are started in hotbeds, soil fumigation is needed to control weeds, soil borne diseases, and insects. Seeds are planted one-quarter to one-half inch deep in rows 4 to 6 inches apart with 2 to 4 seeds per inch and covered with a sash or plastic covering. The seedlings are thinned

at the two-leaf stage allowing 1 ½ inches between plants. Plants are watered twice daily and fertilized with soluble fertilizer at least every 2 weeks. Proper ventilation is important and can be maintained by raising the sash or plastic covering during the hottest portion of the day. In the hotbed, if properly handled, 3 or 4 ounces of seed will produce enough seedlings to plant 1 acre. When seed is planted in beds, it generally requires about 6 to 8 weeks from seed to plants for the spring crop, and about 4 to 5 weeks for the fall crop.

In the greenhouse, a variety of plant growing containers may be used (i.e. plastic cell packs, peat pots, and speedling trays) for growing broccoli. These containers can be filled with an artificial media, usually a combination of peat, perlite, vermiculite, and in some instances bark. The seeds can be sown directly into the containers and thinned upon emergence to 1 plant per cell or pot. In the greenhouse, it generally requires 5 to 6 weeks from seed to plants for the spring crop and 4 to 5 weeks for the fall crop.

For direct seeding, broccoli seeds may be direct seeded in the field using a precision planter. Seed required for one acre is generally 0.75 to 1.25 pound when using a precision seeder.

#### **ORIGIN AND PARENTAGE OF** **HEAT TOLERANT BROCCOLI PLANTS**

The broccoli of this invention were created by classical plant breeding as well as anther culture techniques. The breeding history of the "inbred," "self compatible" and "male" lines identified are exemplified by the following breeding histories.

##### **A. DEVELOPMENT OF INBRED LINES**

Numerous heat tolerant inbred lines were developed. For illustrative, but nonlimiting purposes, the breeding histories of the following inbred lines are presented. Unless otherwise noted, single

plant selections were made each year for plants exhibiting heat tolerance.

1. Inbred Lines 393-2-19, 393-2-47, and 393-2-32

5 All lines designated "393-2-XX" where XX represents a different number for a line were isolated and developed as indicated below. Representative lines include 393-2-19, 393-2-47, 393-2-32.

10 Fall, Year 1 Commercial broccoli hybrid Marathon was crossed with broccoli hybrid "No. 608" obtained from IM Foods, Incorporated, Gilroy, California.

15 Summer, Year 2 F1 seed from Marathon x No. 608 were planted into row number 393 of a summer broccoli trial in Gilroy, California, and single plant selections were made for heat tolerance and downy mildew resistance.

20 Fall, Year 2 Single plant selection number 2 from row 393, i.e. "393-2", which had exhibited good heat tolerance and downy mildew resistance, was entered into anther culture.

25 Spring, Year 3 Anther culture product numbers 19 and 47 from 393-2, i.e. "393-2-19," "393-2-47" and "393-2-32" were transplanted into the greenhouse in Gilroy, California. 393-2-19, 393-2-47 and 393-2-32 were observed to exhibit desirable horticultural traits for deep

30 dome-shaped head, lack of side shoots, good vigor, and high yield. 393-2-19, 393-2-47 and



393-2-32 also demonstrated ability to produce self-pollinated seed.

5 Summer, Year 4 The original seed from 393-2-19, 393-2-47 and 393-2-32 made in the greenhouse in Spring, year 3, were seeded in the greenhouse in Summer, year 4, and subsequently transplanted to the field for evaluation in Gilroy, California, in the summer. 393-2-19 and 393-2-47 exhibited outstanding 10 uniformity and were considered breeding true as a spontaneously doubled-haploid, "inbred", line. Plants were taken from the field plot for self-pollination and subsequent seed increase.

15 From Year 5 to Present 393-2-19, 393-2-47 and 393-2-32 have consistently exhibited exceptionally good uniformity and stability through generations of seed increase with no variants or off types plants 20 ever observed.

25 In the breeding history described above, commercial broccoli hybrid Marathon was crossed with broccoli hybrid No. 608 obtained from IM Foods, Incorporated, Gilroy, California. The commercially available broccoli hybrid Marathon was selected for the initial cross because it had demonstrated good yield potential. Hybrid No. 608 was selected for the initial cross because it was thought to have less side shoots, an advantageous characteristic for harvesting.

30 During the breeding process, F1 seed from Marathon x No. 608 were planted and grown. Selections were made for heat tolerance and downy mildew resistance. The heat tolerance

selection was conducted at head formation through harvest maturity. The selection criteria were smooth domed head, even flower-bud size, good head color, lack of bracting (leaflets in the head), and an ability to hold a good head shape through harvest maturity. The  
5 downy mildew selection was conducted throughout the growth cycle of the plants. The selection for downy mildew resistance was based on plants with no mildew lesions or a greatly reduced number of lesions present on any leaves as compared to non-resistant plants.

Multiple single plant selections exhibiting heat tolerance and  
10 downy mildew resistance were entered into anther culture. Anther culture procedures are well known in the art of plant breeding. In anther culture techniques, the undifferentiated pollen mother cells that exist in immature broccoli anthers are stimulated *in vitro* into embryonic states by procedures well known in the literature. The  
15 undifferentiated pollen mother cells may be subjected to treatments of higher temperatures, light and dark and specialized media growth conditions including hormone simulation. Plant growth conditions of 60°F and bright light followed by a heat shock of 90°F after anther excision and culturing can help stimulate embryogenesis. This  
20 process can stimulate the development of embryonic growth wherein the haploid (one-half the chromosome number) pollen mother cell multiplies and grows into callus tissue. The callus tissue, through the use of specialized media, hormone treatments, and controlled temperature and light can be stimulated to make green plant shoots  
25 and eventually functional roots. Some of these haploids spontaneously double their chromosome number, thus; generating "di-haploids," which are essentially completely homozygous. These highly homozygous lines are genetically equivalent to the end result of many years of self-pollination by conventional means.

30 In a preferred format, anther cultures are prepared as described in Keller, et al., Embryogenesis and Plant Regeneration in

- Brassica napus Anther Cultures, *Canadian Journal of Botany* 55: 1383-1388 (1977); Keller, et al. Production of Haploids via Anther Culture in Brassica Oleracea Var. Italica, *Euphytica* 32: 151-159 (1983); and Orton, et al., Segregation of Genetic Markers among
- 5 Plants Regenerated From Cultured Anthers of Broccoli (Brassica oleracea var. 'italica') *Theor Appl Genet* 69:637-643 (1985).

- In one format, broccoli anthers are prepared and cultured as follows. Immature broccoli racemes are removed from broccoli heads as they begin to elongate, but before the first buds are
- 10 opened. Racemes are then sterilized, for example in 20% W:V household bleach for 15 minutes under agitation with one drop of detergent per 100ml as a surfactant. The racemes are then washed at least three times with sterile distilled water for generally 10 minutes per wash. Anthers are then generally removed by carefully
- 15 peeling back the immature calyx and corolla and gently rupturing the point of filament attached to the anther axis. Care must be taken to minimize damage to the anther culture. Anthers are then placed into liquid culture medium as described in Keller, et al. (1977). Such culture medium may include L-serine at a concentration of 100mg/l.
- 20 Anthers are cultured at, for example, 35°C for 36h and transferred to 25° (all in the dark). In some circumstances, anther-derived embryos are then kept in continuous fluorescent light (25°C) for 1 week to permit greening and then transferred to hormone-free solidified Bs medium as described by Gamborg, et al. *Exp. Cell Res.*
- 25 50: 151-158 (1968). Upon transfer to the hormone-free medium, the anther-derived embryos are cultured at 25°C in light. Embryos which fail to develop into plantlets may be cut into sections and cultured on a modified Murashige and Skoog medium [Murashige, et al., *Physiol. Plant* 15: 473-497 (1962)] containing 0.8% agar, 2%
- 30 sucrose,  $5 \times 10^{-6}$  M benzyladenine and  $10^{-7}$  M naphthaleneacetic acid (NAA) to induce shoot regeneration. To induce rooting, developing

shoots are excised and cultured on B<sub>5</sub> medium in, for example, 60ml sterilin bottles. Rooted plantlets may then be planted in Jiffy-7 peat pellets and kept moist in a mist chamber. After 2 weeks, the broccoli plants can then be transferred to soil and grown in the greenhouse for further selections.


Once transferred to the greenhouse, various plants were selected based on desired phenotypical characteristics including an ability to produce self-pollinated seed. Self-pollination is advantageous because it permits seed increase and bulking of seed without random cross-pollination.

## 2. Inbred lines: 4243, 4221 and 4441

Using the procedures generally outlined above for 393-2-19 and 393-2-47, inbred lines 4243, 4221 and 4441 were isolated following the pedigree chart outlined below. IM Foods 608, Cruiser, Marathon and Sprinter are commercially available lines. Unless otherwise noted, all selections were single plant selections for heat tolerance. 393-2-19 is the same inbred line described above. Each season, the single plant selection exhibiting heat tolerance were selfed and seed was harvested for the next growing season.

20

### Derivation of: 4243; 4221; and 4441



Line 5334-1 X Sprinter F1

(a di-haploid anther  
culture product)



Line 5526-1

↓

Self



Self



Self



Self




Self

1907 1908 1909 1910 1911 1912 1913 1914 1915 1916 1917 1918 1919 1920 1921 1922 1923 1924 1925 1926 1927 1928 1929 1930 1931 1932 1933 1934 1935 1936 1937 1938 1939 1940 1941 1942 1943 1944 1945 1946 1947 1948 1949 1950 1951 1952 1953 1954 1955 1956 1957 1958 1959 1960 1961 1962 1963 1964 1965 1966 1967 1968 1969 1970 1971 1972 1973 1974 1975 1976 1977 1978 1979 1980 1981 1982 1983 1984 1985 1986 1987 1988 1989 1990 1991 1992 1993 1994 1995 1996 1997 1998 1999 2000 2001 2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 2021 2022 2023 2024 2025 2026 2027 2028 2029 2030 2031 2032 2033 2034 2035 2036 2037 2038 2039 2040 2041 2042 2043 2044 2045 2046 2047 2048 2049 2050 2051 2052 2053 2054 2055 2056 2057 2058 2059 2060 2061 2062 2063 2064 2065 2066 2067 2068 2069 2070 2071 2072 2073 2074 2075 2076 2077 2078 2079 2080 2081 2082 2083 2084 2085 2086 2087 2088 2089 2090 2091 2092 2093 2094 2095 2096 2097 2098 2099 2100 2101 2102 2103 2104 2105 2106 2107 2108 2109 2110 2111 2112 2113 2114 2115 2116 2117 2118 2119 2120 2121 2122 2123 2124 2125 2126 2127 2128 2129 2130 2131 2132 2133 2134 2135 2136 2137 2138 2139 2140 2141 2142 2143 2144 2145 2146 2147 2148 2149 2150 2151 2152 2153 2154 2155 2156 2157 2158 2159 2160 2161 2162 2163 2164 2165 2166 2167 2168 2169 2170 2171 2172 2173 2174 2175 2176 2177 2178 2179 2180 2181 2182 2183 2184 2185 2186 2187 2188 2189 2190 2191 2192 2193 2194 2195 2196 2197 2198 2199 2200 2201 2202 2203 2204 2205 2206 2207 2208 2209 2210 2211 2212 2213 2214 2215 2216 2217 2218 2219 2220 2221 2222 2223 2224 2225 2226 2227 2228 2229 2230 2231 2232 2233 2234 2235 2236 2237 2238 2239 2240 2241 2242 2243 2244 2245 2246 2247 2248 2249 2250 2251 2252 2253 2254 2255 2256 2257 2258 2259 2260 2261 2262 2263 2264 2265 2266 2267 2268 2269 2270 2271 2272 2273 2274 2275 2276 2277 2278 2279 2280 2281 2282 2283 2284 2285 2286 2287 2288 2289 2290 2291 2292 2293 2294 2295 2296 2297 2298 2299 2300 2301 2302 2303 2304 2305 2306 2307 2308 2309 2310 2311 2312 2313 2314 2315 2316 2317 2318 2319 2320 2321 2322 2323 2324 2325 2326 2327 2328 2329 2330 2331 2332 2333 2334 2335 2336 2337 2338 2339 2340 2341 2342 2343 2344 2345 2346 2347 2348 2349 2350 2351 2352 2353 2354 2355 2356 2357 2358 2359 2360 2361 2362 2363 2364 2365 2366 2367 2368 2369 2370 2371 2372 2373 2374 2375 2376 2377 2378 2379 2380 2381 2382 2383 2384 2385 2386 2387 2388 2389 2390 2391 2392 2393 2394 2395 2396 2397 2398 2399 2400 2401 2402 2403 2404 2405 2406 2407 2408 2409 2410 2411 2412 2413 2414 2415 2416 2417 2418 2419 2420 2421 2422 2423 2424 2425 2426 2427 2428 2429 2430 2431 2432 2433 2434 2435 2436 2437 2438 2439 2440 2441 2442 2443 2444 2445 2446 2447 2448 2449 2450 2451 2452 2453 2454 2455 2456 2457 2458 2459 2460 2461 2462 2463 2464 2465 2466 2467 2468 2469 2470 2471 2472 2473 2474 2475 2476 2477 2478 2479 2480 2481 2482 2483 2484 2485 2486 2487 2488 2489 2490 2491 2492 2493 2494 2495 2496 2497 2498 2499 2500 2501 2502 2503 2504 2505 2506 2507 2508 2509 2510 2511 2512 2513 2514 2515 2516 2517 2518 2519 2520 2521 2522 2523 2524 2525 2526 2527 2528 2529 2530 2531 2532 2533 2534 2535 2536 2537 2538 2539 2540 2541 2542 2543 2544 2545 2546 2547 2548 2549 2550 2551 2552 2553 2554 2555 2556 2557 2558 2559 2560 2561 2562 2563 2564 2565 2566 2567 2568 2569 2570 2571 2572 2573 2574 2575 2576 2577 2578 2579 2580 2581 2582 2583 2584 2585 2586 2587 2588 2589 2590 2591 2592 2593 2594 2595 2596 2597 2598 2599 2600 2601 2602 2603 2604 2605 2606 2607 2608 2609 2610 2611 2612 2613 2614 2615 2616 2617 2618 2619 2620 2621 2622 2623 2624 2625 2626 2627 2628 2629 2630 2631 2632 2633 2634 2635 2636 2637 2638 2639 2640 2641 2642 2643 2644 2645 2646 2647 2648 2649 2650 2651 2652 2653 2654 2655 2656 2657 2658 2659 2660 2661 2662 2663 2664 2665 2666 2667 2668 2669 2670 2671 2672 2673 2674 2675 2676 2677 2678 2679 2680 2681 2682 2683 2684 2685 2686 2687 2688 2689 2690 2691 2692 2693 2694 2695 2696 2697 2698 2699 2700 2701 2702 2703 2704 2705 2706 2707 2708 2709 2710 2711 2712 2713 2714 2715 2716 2717 2718 2719 2720 2721 2722 2723 2724 2

3. Inbred lines: 4274-1; 4274-2; 4278-1; 4284-1; 4285-1;  
4354-1; 4354-2; 4377-1


Inbred lines 4274-1; 4274-2; 4278-1; 4284-1; 4285-1; 4354-1; 4354-2; 4377-1 were isolated following the pedigree chart outlined below using the procedures generally described above for inbred lines 393-2-19 and 393-2-47. Unless otherwise noted, single plant selections were made for heat tolerance.



4. Inbred lines: 4318-1; 4320-1 and 4320-2; 4321-1; 4437-1; 4476-1; 4462-1

Inbred lines 4318-1; 4320-1 and 4320-2; 4321-1; 4437-1; 4476-1; 4462-1 were isolated following the pedigree chart outlined below using the procedures generally outlined above for inbred lines 393-2-19 and 393-2-47. Commercially available line Marathon was selfed. Repeated single plant selections for heat tolerance were made to produce the resulting lines. In year 6, the selected line was mass pollinated.

Derivation of: 4318-1; 4320-1 and 4320-2; 4321-1; 4437-1; 4476-1; 4462-1




5. Inbred lines: 4308-2; 4309-1; 4355-1; 4412-1; 4301; 4303;  
4304; 4317; 4468; 4470; 4471

Inbred lines 4308-2; 4309-1; 4355-1; 4412-1; 4301; 4303;  
 4304; 4317; 4468; 4470; 4471 were isolated following the

- 5 pedigree chart outlined below using the techniques generally outlined  
 above for inbred lines 393-2-19 and 393-2-47. In year 9, the  
 selected lines were brush pollinated, i.e., pollinated with a brush.  
 Single plant selections were made for heat tolerance.

Derivation of: 4308-2; 4309-1; 4355-1; 4412-1;  
 10 4301; 4303; 4304; 4317; 4468; 4470; 4471






6. Inbred lines 4263-1; 4430-1; 4450-1 and 4450-2:

Inbred lines 4263-1; 4430-1; 4450-1 and 4450-2 were isolated following the pedigree chart outlined below using the techniques generally outlined above for inbred lines 393-2-19 and 393-2-47. Single plant selections were made for heat tolerance.


Derivation of: 4263-1; 4430-1; 4450-1; 4450-2



# 7. Inbred line 4432-1

Inbred line 4432-1 was isolated following the pedigree chart outlined below using the techniques generally outlined above for inbred lines 393-2-19 and 393-2-47. Single plant selections for heat tolerance were made.


## Derivation of: 4432-1



8. Inbred line 4267-1 ( = "2192")

Inbred line 4267-1 ( = "2192") was isolated following the pedigree chart outlined below using the techniques generally outlined above for inbred line 393-2-19. Single plant selections were made  
5 for heat tolerance.


Derivation of: 4267-1 ( = "2192")



9. Inbred line 7861


Inbred line 7861 was isolated following the pedigree chart outlined below using the techniques generally outlined for inbred lines 393-2-19 and 393-2-47 outlined above. Single plant selections were made for heat tolerance.

Derivation of: 7861



**10. Inbred line 7864**


Inbred line 7864 was isolated following the pedigree chart outlined below using the techniques generally outlined for inbred lines 393-2-19 and 393-2-47 outlined above. Single plant selections were made for heat tolerance.



**11. Inbred line 7865**

Inbred line 7865 was isolated following the pedigree chart outlined below using the techniques generally outlined for inbred lines 393-2-19 and 393-2-47 outlined above. Single plant selections were made for heat tolerance.

**Derivation of: 7865**



1. 1971-1972	1973-1974	1975-1976	1977-1978	1979-1980	1981-1982	1983-1984	1985-1986	1987-1988	1989-1990	1991-1992	1993-1994	1995-1996	1997-1998	1999-2000	2001-2002	2003-2004	2005-2006	2007-2008	2009-2010	2011-2012	2013-2014	2015-2016	2017-2018	2019-2020	2021-2022	2023-2024	2025-2026	2027-2028	2029-2030	2031-2032	2033-2034	2035-2036	2037-2038	2039-2040	2041-2042	2043-2044	2045-2046	2047-2048	2049-2050	2051-2052	2053-2054	2055-2056	2057-2058	2059-2060	2061-2062	2063-2064	2065-2066	2067-2068	2069-2070	2071-2072	2073-2074	2075-2076	2077-2078	2079-2080	2081-2082	2083-2084	2085-2086	2087-2088	2089-2090	2091-2092	2093-2094	2095-2096	2097-2098	2099-2100	2101-2102	2103-2104	2105-2106	2107-2108	2109-2110	2111-2112	2113-2114	2115-2116	2117-2118	2119-2120	2121-2122	2123-2124	2125-2126	2127-2128	2129-2130	2131-2132	2133-2134	2135-2136	2137-2138	2139-2140	2141-2142	2143-2144	2145-2146	2147-2148	2149-2150	2151-2152	2153-2154	2155-2156	2157-2158	2159-2160	2161-2162	2163-2164	2165-2166	2167-2168	2169-2170	2171-2172	2173-2174	2175-2176	2177-2178	2179-2180	2181-2182	2183-2184	2185-2186	2187-2188	2189-2190	2191-2192	2193-2194	2195-2196	2197-2198	2199-2200	2201-2202	2203-2204	2205-2206	2207-2208	2209-2210	2211-2212	2213-2214	2215-2216	2217-2218	2219-2220	2221-2222	2223-2224	2225-2226	2227-2228	2229-2230	2231-2232	2233-2234	2235-2236	2237-2238	2239-2240	2241-2242	2243-2244	2245-2246	2247-2248	2249-2250	2251-2252	2253-2254	2255-2256	2257-2258	2259-2260	2261-2262	2263-2264	2265-2266	2267-2268	2269-2270	2271-2272	2273-2274	2275-2276	2277-2278	2279-2280	2281-2282	2283-2284	2285-2286	2287-2288	2289-2290	2291-2292	2293-2294	2295-2296	2297-2298	2299-2300	2301-2302	2303-2304	2305-2306	2307-2308	2309-2310	2311-2312	2313-2314	2315-2316	2317-2318	2319-2320	2321-2322	2323-2324	2325-2326	2327-2328	2329-2330	2331-2332	2333-2334	2335-2336	2337-2338	2339-2340	2341-2342	2343-2344	2345-2346	2347-2348	2349-2350	2351-2352	2353-2354	2355-2356	2357-2358	2359-2360	2361-2362	2363-2364	2365-2366	2367-2368	2369-2370	2371-2372	2373-2374	2375-2376	2377-2378	2379-2380	2381-2382	2383-2384	2385-2386	2387-2388	2389-2390	2391-2392	2393-2394	2395-2396	2397-2398	2399-2400	2401-2402	2403-2404	2405-2406	2407-2408	2409-2410	2411-2412	2413-2414	2415-2416	2417-2418	2419-2420	2421-2422	2423-2424	2425-2426	2427-2428	2429-2430	2431-2432	2433-2434	2435-2436	2437-2438	2439-2440	2441-2442	2443-2444	2445-2446	2447-2448	2449-2450	2451-2452	2453-2454	2455-2456	2457-2458	2459-2460	2461-2462	2463-2464	2465-2466	2467-2468	2469-2470	2471-2472	2473-2474	2475-2476	2477-2478	2479-2480	2481-2482	2483-2484	2485-2486	2487-2488	2489-2490	2491-2492	2493-2494	2495-2496	2497-2498	2499-2500	2501-2502	2503-2504	2505-2506	2507-2508	2509-2510	2511-2512	2513-2514	25
--------------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	-----------	----


	Year 1	IMF 608	X	Green Belt
			↓	
10	Year 2		X	Green Belt
			↓	
	Year 3		Self	
			↓	
	Year 4		Self	
15			↓	
	Year 5		Self	
			↓	
	Year 6		Self	
			↓	
20	Year 7		Self	
			↓	
	Year 8		Self	
			↓	
	Year 9		Self	





**14. Inbred line 7935**

Inbred line 7935 was isolated following the pedigree chart outlined below using the techniques generally outlined for inbred lines 393-2-19 and 393-2-47 outlined above. Single plant selections were made for heat tolerance.



# 15. Inbred line 8092

Inbred line 8092 was isolated following the pedigree chart outlined below using the techniques generally outlined for inbred lines 393-2-19 and 393-2-47 outlined above. Single plant selections were made for heat tolerance.

## Derivation of: 8092


	Year 1	Cruiser	X	Green Belt
			↓	
10	Year 2		X	Marathon
			↓	
	Year 3		Self	
			↓	
	Year 4		Self	
15			↓	
	Year 5		Self	
			↓	
	Year 6		Self	
			↓	
20	Year 7		Self	
			↓	
	Year 8		Self	
			↓	
	Year 9		Self	

1. The first of these is the fact that the
 2. second of these is the fact that the
 3. third of these is the fact that the
 4. fourth of these is the fact that the
 5. fifth of these is the fact that the
 6. sixth of these is the fact that the
 7. seventh of these is the fact that the
 8. eighth of these is the fact that the
 9. ninth of these is the fact that the
 10. tenth of these is the fact that the
 11. eleventh of these is the fact that the
 12. twelfth of these is the fact that the
 13. thirteenth of these is the fact that the
 14. fourteenth of these is the fact that the
 15. fifteenth of these is the fact that the
 16. sixteenth of these is the fact that the
 17. seventeenth of these is the fact that the
 18. eighteenth of these is the fact that the
 19. nineteenth of these is the fact that the
 20. twentieth of these is the fact that the
 21. twenty-first of these is the fact that the
 22. twenty-second of these is the fact that the
 23. twenty-third of these is the fact that the
 24. twenty-fourth of these is the fact that the
 25. twenty-fifth of these is the fact that the
 26. twenty-sixth of these is the fact that the
 27. twenty-seventh of these is the fact that the
 28. twenty-eighth of these is the fact that the
 29. twenty-ninth of these is the fact that the
 30. thirtieth of these is the fact that the
 31. thirty-first of these is the fact that the
 32. thirty-second of these is the fact that the
 33. thirty-third of these is the fact that the
 34. thirty-fourth of these is the fact that the
 35. thirty-fifth of these is the fact that the
 36. thirty-sixth of these is the fact that the
 37. thirty-seventh of these is the fact that the
 38. thirty-eighth of these is the fact that the
 39. thirty-ninth of these is the fact that the
 40. fortieth of these is the fact that the
 41. forty-first of these is the fact that the
 42. forty-second of these is the fact that the
 43. forty-third of these is the fact that the
 44. forty-fourth of these is the fact that the
 45. forty-fifth of these is the fact that the
 46. forty-sixth of these is the fact that the
 47. forty-seventh of these is the fact that the
 48. forty-eighth of these is the fact that the
 49. forty-ninth of these is the fact that the
 50. fiftieth of these is the fact that the
 51. fifty-first of these is the fact that the
 52. fifty-second of these is the fact that the
 53. fifty-third of these is the fact that the
 54. fifty-fourth of these is the fact that the
 55. fifty-fifth of these is the fact that the
 56. fifty-sixth of these is the fact that the
 57. fifty-seventh of these is the fact that the
 58. fifty-eighth of these is the fact that the
 59. fifty-ninth of these is the fact that the
 60. sixtieth of these is the fact that the
 61. sixty-first of these is the fact that the
 62. sixty-second of these is the fact that the
 63. sixty-third of these is the fact that the
 64. sixty-fourth of these is the fact that the
 65. sixty-fifth of these is the fact that the
 66. sixty-sixth of these is the fact that the
 67. sixty-seventh of these is the fact that the
 68. sixty-eighth of these is the fact that the
 69. sixty-ninth of these is the fact that the
 70. seventieth of these is the fact that the
 71. seventy-first of these is the fact that the
 72. seventy-second of these is the fact that the
 73. seventy-third of these is the fact that the
 74. seventy-fourth of these is the fact that the
 75. seventy-fifth of these is the fact that the
 76. seventy-sixth of these is the fact that the
 77. seventy-seventh of these is the fact that the
 78. seventy-eighth of these is the fact that the
 79. seventy-ninth of these is the fact that the
 80. eightieth of these is the fact that the
 81. eighty-first of these is the fact that the
 82. eighty-second of these is the fact that the
 83. eighty-third of these is the fact that the
 84. eighty-fourth of these is the fact that the
 85. eighty-fifth of these is the fact that the
 86. eighty-sixth of these is the fact that the
 87. eighty-seventh of these is the fact that the
 88. eighty-eighth of these is the fact that the
 89. eighty-ninth of these is the fact that the
 90. ninetieth of these is the fact that the
 91. ninety-first of these is the fact that the
 92. ninety-second of these is the fact that the
 93. ninety-third of these is the fact that the
 94. ninety-fourth of these is the fact that the
 95. ninety-fifth of these is the fact that the
 96. ninety-sixth of these is the fact that the
 97. ninety-seventh of these is the fact that the
 98. ninety-eighth of these is the fact that the
 99. ninety-ninth of these is the fact that the
 100. hundredth of these is the fact that the

10


15

## IMF 608 Self



**17. Inbred line 7914**


Inbred line 7914 was isolated following the pedigree chart outlined below using the techniques generally outlined for inbred lines 393-2-19 and 393-2-47 outlined above. Single plant selections  
5 were made for heat tolerance.



**18. Inbred lines 7770 and 5580-2**

Inbred lines 7770 and 5580-2 were isolated following the pedigree chart outlined below using the techniques generally outlined for inbred lines 393-2-19 and 393-2-47 outlined above. Single plant  
5 selections were made for heat tolerance.


**Derivation of: 7770 and 5580-2**



**19. Inbred line 7778**

Inbred line 7778 was isolated following the pedigree chart outlined below using the techniques generally outlined for inbred lines 393-2-19 and 393-2-47 outlined above. Single plant selections were made for heat tolerance.

**Derivation of: 7778**



## B. SELF INCOMPATIBLE LINES


Numerous heat tolerant self-incompatible ("female") lines were developed. For illustrative, but not limiting purposes, the breeding histories of the following self-incompatible lines are presented.

5. Unless otherwise noted, single plant selections were made for heat tolerance.

### 1. Self-incompatible lines: 4201; 4219, 4237, 4280, 4287, 4288, 4289, 4290, 4291, 4458-1, 4460-1


- 10 Broccoli lines 4201; 4219, 4237, 4280, 4287, 4288, 4289, 4290, 4291, 4458-1, 4460-1 were isolated following the pedigree chart outlined below using the procedures generally outlined above for the isolation of 393-2-19.

### Derivation of: 4201; 4219, 4237, 4280, 4287, 4288, 4289, 4290, 4291, 4458-1, 4460-1



2. Self Incompatible Line: 4415

Broccoli line 4415 was isolated following the pedigree chart outlined below using the procedures generally outlined above for 393-2-19.






3. Self Incompatible Line: 4418


Broccoli line 4418 was isolated following the pedigree chart outlined below using the procedures outlined above for 393-2-19.

Derivation of: 4418



4. Self Incompatible Line 4395-2

Broccoli line 4395-2 was isolated following the pedigree chart outlined below using the procedures generally outlined above for 4935-2.



**C. MALE LINES**

Numerous heat tolerant "male" broccoli lines have been identified and shown stable and uniform. For illustrative but non-limiting purposes, the breeding histories of the M7007, M7009 and  
5 M7028 are provided as follows.

The "Cruiser" broccoli line was selected for initial crosses because it was a commercially available hybrid that showed a small degree of heat tolerance which was rated at approximately 5 and also had a nicely elevated head.

10

**M7007**

Year One IM Hybrid No. 608 obtained from IM Foods, Incorporated, Gilroy, California, was self-pollinated.

Year Two F2 of Hybrid No. 608 is crossed with Cruiser, which  
15 was obtained from Royal Sluis, a Dutch seed company.

Year Three Heat tolerant single plant selection of the F2 Hybrid 608/Cruiser with heat tolerance equaling 8- was made.

Year Four Heat tolerance equaling 7 + single plant selection gives [(No. 608) F2/Cruiser] F3.

20 Year Five Eight selections are selfed and massed selected to give [(No. 608) F2/Cruiser] F4.

Year Six Twelve selections are massed.

Year Seven Fifteen selections are massed.

Year Eight Five selections are massed selected and entered into a  
25 large isolation cage increase to give the finished line M7007.

**M7009**

- Year One IM Hybrid No. 608 is self-pollinated.
- Year Two F2 of Hybrid No. 608 is crossed with Cruiser.
- Year Three Heat tolerant single plant selection of the F2 Hybrid
- 5 608/Cruiser with heat tolerance equaling 8- was made.
- Year Four Heat tolerance equaling 7 + single plant selection gives [(No. 608) F2/Cruiser] F3.
- Year Five Eight selections are selfed and massed to give [(No. 608) F2/Cruiser] F4.
- 10 Year Six Twelve selections are massed.
- Year Seven Fifteen selections are massed.
- Year Eight Five selections are massed selected and entered into a large isolation cage increase to give the finished line.
- Year Nine Seed storage.
- 15 Year Ten Six selections are massed selected and entered into a large isolation cage increase to give finish line M7009.

**M7028**

- Year Two F2 of Hybrid No. 608 is crossed with Cruiser.
- 20 Year Three Heat tolerant single plant selection of the F2 Hybrid 608/Cruiser with heat tolerance equaling 8- was made.
- Year Four Made single plant selection. [(No. 608) F2/Cruiser] F3
- Year Five Made single plant selection with heat tolerance equaling 7. [(No. 608) F2/Cruiser] F4
- 25 Year Six Made single plant selection with heat tolerance equaling 7. [(No. 608) F2/Cruiser] F5
- Year Seven Selected five plants, massed selected and entered into a large isolation cage to give finished line M7028.
- 30 The male lines of this invention can be crossed with female lines (self-incompatible) to produce hybrid seed. The female lines

may or may not be heat tolerant. Encompassed within the scope of this invention are the hybrid seed produced from crossing the male lines of this invention with other broccoli lines of interest. Hybrid seed includes but is not limited to H7007, H7008, H7028.

5

#### **HYBRID SEED PRODUCTION**

For hybrid seed production of heat tolerant broccoli seed, two lines are selected for production. The lines are designated male or female, with the female being the recipient of the male line pollen.

- 10 Either the male or female or both lines may be heat tolerant as defined by this invention. Broccoli plants flower with both the female and male parts and are capable of self-pollination. The line designated "female" is generally "self-incompatible," which means it will not accept its own pollen, a process developed in the plant by
- 15 breeding. The line designated "male" is generally "self-compatible" and will accept its own pollen. Since self-incompatible lines will not accept their own pollen, but will out-cross with other broccoli pollen. Self-incompatible lines produce the commercially desired hybrid seed. The male line is the pollen provider to the female line. The
- 20 cross of the self-compatible male line and the self-incompatible female line will produce a seed that is a hybrid.

- Once a hybrid has been selected for seed production, a "nick" study is done. The nick study identifies the flowering period of the female, i.e. when it will start to flower and for how long it will
- 25 flower. A nick study is also done for the male line and the two are compared. The nick study gives the data needed to determine if the female will require 1, 2, 3, or 4 male planting dates to cover its full flowering period.

- Once the data from the nick study is obtained, seed of the
- 30 female and the first male are planted in the greenhouse. The second male is planted in the greenhouse 7 - 10 days later, with the third

male planted another 7 – 10 days after that, and the final or 4<sup>th</sup> male planted within 10 more days. The female is seeded in the greenhouse at approximately 11,000 plants for each production acre and each male planting at 7,000 plants. Forty-five days from being  
5 planted in the greenhouse the female and first male are ready for transplanting in the field. The three remaining males are each transplanted into the field within 45 days of their individual greenhouse planting dates.

#### 10 Field Production of Hybrid Seed

Field production of the hybrid seed is begun when all of the female plants and the first male plants from the greenhouse are transplanted into the field. Transplanting can be done by machine or by hand with large crews. The plants are placed into the soil on  
15 prepared listed beds that are on 40-inch centers (see Figure 2). The depth of the planting is generally 3 inches, but depends on the size of the transplant plug. Each plant is separated approximately eighteen inches apart going down the seed-line and each parallel seed-line on a single bed is twelve inches apart. The successive  
20 plantings of the second, third, and fourth male follow the female planting at approximately ten day intervals. An illustrative planting schedule is as follows:

	October 15	female transplanting date
	October 15	first male transplanting date
25	October 25	second male transplanting date
	November 4	third male transplanting date
	November 14	fourth male transplanting date

The dates are not fixed, but are an approximation for illustrative and non-limiting purposes.

30 Once all the plantings are accomplished, the field is watched for typical cultural problems found in all broccoli production, whether

for seed or vegetable. These problems include weeds, diseases, insect pests, irrigation, fertilization, and cultivation.

The singular difference for a seed production field as compared to a broccoli production field is the use of roguing.

- 5 Roguing is simply the walking through and examination of a field and checking each plant for correctness to type. Any plant that does not fit the proper description for type is pulled and destroyed or "rogued." The roguing starts within thirty days of the last male transplanted and continues until the field is at a market ready point,
- 10 which is generally 100 days. Once the field is at market ready (market ready being the point where the heads are harvestable as a vegetable for sale) the seed production starts. Market ready heads are generally seen in the female and first male in late April to early May of the year following transplanting. The fully developed heads
- 15 age and then bolt, which is the extension of individual flower stalks. The nick or timing between the male plants bolting and female plants bolting is now the crucial item watched. The female will only set hybrid seed if pollen is in constant and abundant supply from a male plant. The heads of both the male and female plant can be trimmed
- 20 to accelerate or slow down the flowering to insure abundant male flowers are available as the female plant flowers. Pollen transfer from the male to the female is done by honeybees, which are commercially supplied. Each acre of seed production requires three to five hives of honeybees. The flowering stage will last sixty to
- 25 eighty days.

- The flowering period is followed by the maturation of the seed within seedpods. The maturation period of 40 to 60 days is checked by monitoring the seed development, as it goes from green and water filled, to the dough stage ending with the seed turning from
- 30 green to brown in color. A judgment call is made, measuring the number of the mature seeds versus seeds yet completed. When the

majority of the seed is mature the female plants are cut by hand and laid in rows (windrowed) to dry down for combining. Ten to twenty days are needed for the plants to dry down.

Combining is a process, which entails the use of a large  
5 harvest machine that lifts the broccoli plants from the ground and grinds them for seed preparation. The plant material is cleaned away from the seed by screens and air, leaving only seed. Combining is the initiation of the seed re-conditioning process. Once combined or harvested, the seed is sent to a mill, which further cleans the seed,  
10 separates the clean seed by size and weight within a size. All testing for purity, disease, germination, and percent hybridity is done on the clean, sized, and weighted seed. If the seed passes the testing it is canned for sale.

The above method describes the seed production methods for  
15 the specific hybrids H7007, H7009, H7022, H2061, H2088, H7021R and H7028 and generally is the method used for all other hybrid broccoli seed production. Hybrid seeds H7007, H7009 H7022, H2061, H2088, H7021R and H70028 were produced by crossing corresponding male lines with 393-2-19 as follows:

- 20 H7007 = 393-2-19 X male 7007  
H7009 = 393-2-19 X male 7009  
H7022 = 393-2-19 X male 5580-2 (same derivation as 7770)  
H2061 = 393-2-19 X male 1551 (same derivation as 7881)  
H2088 = 393-2-19 male 7009  
25 H7021R = 5580 (same derivation as 7770) X 393-2-19  
H7028 = 393-2-19 X male 7028

#### COMPARATIVE STUDIES

Several studies have been performed to compare and contrast  
30 the broccoli lines of this invention with commercially available broccoli lines.



**Comparative Analysis Study #1**

In study #1, the following broccoli lines were analyzed: Hybrid 7007, hybrid 7008, hybrid 7022, hybrid 7028, male 7007, male 7009, male 7022, male 7028, hybrid 393-2-19, hybrid 393-2-47, 5 Marathon and Pinnacle. Marathon and Pinnacle are commercially available broccoli hybrids. Hybrid 7022 resulted from a cross between 393-12-19 and 5580-2 (393-2-19/5580-2). As indicated above, 5580 is the same derivative as 7770.

Broccoli seeds were sown in the greenhouse. Broccoli 10 seedlings were transplanted to the field on August 8. Daily high and low temperature measurements during the course of study #1 are presented in Table 1. Note that the growing temperatures for study #1 were generally quite warm.

In study #1, the days from direct seeding to 50% harvest; 15 days from transplanting to 50% harvest and the length of the harvest period are shown in Table 2.1. The results indicate that the broccoli lines of this invention have a significantly longer harvest period than the commercially available hybrids Marathon and Pinnacle. A longer period in which the head remains available for 20 harvest offers growers greater flexibility in harvesting and therefore greatly reduces costs. The harvest "holding" ability is due, in part, to heat tolerance.

Table 2.2 shows data summarizing various characteristics of the broccoli plants at harvest. Tables 2.3A and 2.3B show data 25 regarding the characteristics of the outer leaves at harvest. The data indicate that both Pinnacle and Marathon were gray-green in foliage color, which is demonstratively different and less commercially acceptable than the blue green foliage of the heat tolerant lines of the invention.

30 Table 2.4A-2.4D show characteristics of the broccoli heads at market maturity. Table 2.5 shows flower color.

5

### Temperature Data for Study #1

Date	Temperature (°F)		
	Max	min	Average
07/03	79	47	62
07/04	84	48	63
07/05	75	52	60
07/06	75	52	62
07/07	75	52	61
07/08	71	48	59
07/09	63	54	58
07/10	70	54	59
07/11	73	53	59
07/12	74	52	59
07/13	85	53	65
07/14	87	54	69
07/15	82	50	63
07/16	72	32	62
07/17	76	56	64
07/18	83	58	68
07/19	89	52	69
07/20	83	53	67
07/21	88	53	71
07/22	100	55	78
07/23	99	59	77
07/24	88	56	69
07/25	95	54	72
07/26	81	58	70
07/27	76	55	63
07/28	78	55	62
07/29	75	56	62
07/30	72	56	61
07/31	72	57	62
08/01	82	57	65
08/02	83	56	65
08/03	88	54	68
08/04	83	56	66
08/05	77	56	64
08/06	74	58	64
08/07	79	59	66
08/08	90	56	71
08/09	98	59	74
08/10	109	60	81
08/11	100	61	78
08/12	91	58	70

### Temperature Data for Study #1

Date	Temperature (°F)		
	Max	min	Average
08/13	83	56	66
08/14	84	52	64
08/15	81	50	62
08/16	86	50	66
08/17	92	53	71
08/18	98	58	75
08/19	97	60	75
08/20	92	57	71
08/21	89	58	68
08/22	74	54	61
08/23	74	53	61
08/24	72	51	61
08/25	69	53	60
08/26	71	52	60
08/27	77	54	63
08/28	81	50	63
08/29	86	51	67
08/30	87	50	67
08/31	83	51	64
09/01	69	54	59
09/02	81	51	61
09/03	91	48	64
09/04	95	52	71
09/05	94	46	72
09/06	95	57	71
09/07	91	55	69
09/08	93	54	71
09/09	98	54	69
09/10	91	58	71
09/11	83	61	69
09/12	89	64	73
09/13	95	68	77
09/14	93	64	75
09/15	77	56	64
09/16	68	56	60
09/17	83	32	70
09/18	85	52	67
09/19	87	55	69
09/20	88	57	68
09/21	76	55	62
09/22	73	53	61

**Table 2: Comparative Analysis (Study #1)**

**2.1. Region of Adaption/Maturity Main Crop at 50% Harvest**

#	I.D.	Region of Adaption	Days from Direct Seeding to 50% Harvest	Days from Transplanting to 50% Harvest	Length of Harvest Period in Days
1	Hybrid 7007	Most regions	137	84	7
2	Hybrid 7008	Most regions	137	87	6
3	Hybrid 7022	Most regions	127	77	6
4	Hybrid 7028	Most regions	136	86	6
5	Male 7007	Southwest	135	85	3
6	Male 7009	Southwest	135	85	4
7	Male 7022	Southwest	123	73	5
8	Male 7028	Southwest	138	88	5
9	Hybrid 393-2-19	Most regions	137	87	8
10	Hybrid 393-2-47	Most regions	133	83	6
11	Marathon	Most regions	134	84	4
12	Pinnacle	Southwest	123	73	2

2.2. Study #1  
Plant (At Harvest)

#	I.D.	Plant Height (cm)	Head Height (cm)	Plant Branches	Plant Habit	Market Class	Lifecycle	Variety Type
1	Hybrid 7007	76.5	57.5	Few	Intermediate	Fresh Market/ Processing	Annual	First generation hybrid
2	Hybrid 7008	-	-	-	-	-	-	-
3	Hybrid 7022	72	51.5	Few	Spreading	Fresh Market/ Processing	Annual	First generation hybrid
4	Hybrid 7028	82.5	57.5	Few	Intermediate	Fresh Market/ Processing	Annual	First generation hybrid
5	Male 7007	92	65	Few	Intermediate	Fresh Market/ Processing	Annual	Inbred
6	Male 7009	92	76	Few	Intermediate	Fresh Market/ Processing	Annual	Inbred
7	Male 7022	58	35	Few	Compact	Fresh Market/ Processing	Annual	Inbred
8	Male 7028	74.5	51	Few	Intermediate	Fresh Market/ Processing	Annual	Inbred

2.2. Study #1  
Plant (At Harvest)

#	I.D.	Plant Height (cm)	Head Height (cm)	Plant Branches	Plant Habit	Market Class	Lifecycle	Variety Type
9	Inbred 393-2-19	62	45.5	Few	Intermediate	Fresh Market/ Processing	Annual	Inbred
10	Inbred 393-2-47	60	48.5	Few	Intermediate	Fresh Market/ Processing	Annual	Inbred
11	Marathon	86.5	56.5	Medium	Spreading	Fresh Market/ Processing	Annual	First generation hybrid
12	Pinnacle	88.5	61.5	Few	Intermediate	Fresh Market/ Processing	Annual	First generation hybrid

702290 2006 9860

2.3A. Study #1  
Outer Leaves (At Harvest)

#	I.D.	# Leaves Per Plant	Leaf Width (cm)	Leaf Length (cm)	Petiole Length (cm)	Leaf Attachment	Wax Presence	Foliage Color
1	Hybrid 7007	29	20	52.5	22	Petiolate	Strong	Blue-green
2	Hybrid 7008	-	-	-	-	-	-	-
3	Hybrid 7022	18	14	41	17.5	Petiolate	Strong	Blue-green
4	Hybrid 7028	25	17.5	53.5	19.5	Petiolate	Strong	Blue-green
5	Male 7007	30	16.5	40.5	13.5	Petiolate	Strong	Blue-green
6	Male 7009	26	15.5	47	18	Petiolate	Strong	Blue-green
7	Male 7022	21	23.5	48	17.5	Petiolate	Strong	Blue-green
8	Male 7028	34	15.5	42.5	19.	Petiolate	Strong	Blue-green
9	Inbred 393- 2-19	23	14	36	11.5	Petiolate	Strong	Blue-green
10	Inbred 393- 2-47	24	16.5	40	17.5	Petiolate	Strong	Blue-green
11	Marathon	50	15.5	50	22	Petiolate	Intermediate	Grey-green
12	Pinnacle	27	16	46.5	21	Petiolate	Intermediate	Grey-green



T02290" 20069360

2.3B. Study #1  
Outer Leaves (At Harvest)

#	I.D.	Leaf Shape	Leaf Base	Leaf Apex	Leaf Margins	Leaf Veins	Midrib	Blistering	Attitude	Leaf tip Torsion	Upper Side of Leaf Profile
1	Hybrid 7007	Elliptic	Blunt	Blunt	Slightly wavy	Inter-mediate	Slightly raised	None	Semi-erect	Weak	Concave
2	Hybrid 7008	-	-	-	-	-	-	-	-	-	-
3	Hybrid 7022	Narrow elliptic	Blunt	Blunt	Slightly wavy	Inter-mediate	Slightly raised	None	Semi-erect	None	Planar
4	Hybrid 7028	Elliptic	Blunt	Blunt	Slightly wavy	Inter-mediate	Slightly raised	None	Semi-erect	None	Concave
5	Male 7007	Elliptic	Blunt	Blunt	Slightly wavy	Inter-mediate	Slightly raised	None	Erect	Weak	Concave
6	Male 7009	Elliptic	Pointed	Blunt	Slightly wavy	Inter-mediate	Slightly raised	None	Erect	Intermediate	Planar
7	Male 7022	Broad elliptic	Blunt	Blunt	Slightly wavy	Inter-mediate	Not raised	None	Horizontal/Semi-erect	Weak	Concave

2.3B. Study #1  
Outer Leaves (At Harvest)

#	I.D.	Leaf Shape	Leaf Base	Leaf Apex	Leaf Margins	Leaf Veins	Midrib	Blistering	Attitude	Leaf tip Torsion	Upper Side of Leaf Profile
8	Male 7028	Elliptic	Blunt	Blunt	Slightly wavy	Inter-mediate	Slightly raised	None	Erect	None	Concave
9	Inbred 393-2-19	Elliptic	Blunt	Blunt	Slightly wavy	Inter-mediate	Slightly raised	None	Erect	None	Concave
10	Inbred 393-2-47	Elliptic	Blunt	Blunt	Slightly wavy	Inter-mediate	Slightly raised	None	Semi-erect	None	Concave
11	Marathon	Narrow elliptic	Blunt	Blunt	Very wavy	Inter-mediate	Slightly raised	None	Horizontal/ Semi-erect	None	Concave
12	Pinnacle	Elliptic	Blunt	Blunt	Slightly wavy	Inter-mediate	Slightly raised	None	Horizontal	None	Concave

**2.4A. Study #1**  
**Head (At Market Maturity)**

#	I.D.	Head Diameter (cm)	Head Depth (cm)	Head Weight (gm)	Head Color	Head Shape
1	Hybrid 7007	17	13.5	446.3	Blue/Green	Transverse broad elliptic
2	Hybrid 7008	-	-	352	-	-
3	Hybrid 7022	15	10.5	377.2	Blue/Green	Transverse narrow elliptic
4	Hybrid 7028	15	11.5	364.4	Blue/Green	Transverse broad elliptic
5	Male 7007	10	8	93.7	Blue/Green	Circular
6	Male 7009	10	8.5	126.2	Blue/Green	Transverse broad elliptic
7	Male 7022	15	10	289.5	Blue/Green	Transverse narrow elliptic
8	Male 7028	11	9.5	165.9	Blue/Green	Circular
9	Inbred 393- 2-19	13	10	325.7	Blue/Green	Transverse elliptic
10	Inbred 393- 2-47	11.5	9	194.7	Blue/purple	Transverse broad elliptic
11	Marathon	14.5	12.5	300	Medium green	Transverse elliptic
12	Pinnacle	14	9.5	274.6	Medium green	Transverse elliptic
13	Male 7008	-	-	150	-	-

702230 20063260

2.4B. Study #1  
Head (At Market Maturity)

#	I.D.	Dome Shape	Head Size	Compactness	Surface Knobbling	Bead Size	Flower Buds
1	Hybrid 7007	Semi-domed	Large	Short pedicels (tight)	Fine	Medium	Even in size
2	Hybrid 7008	-	-	-	-	-	-
3	Hybrid 7022	Very Deeply domed	Large	Short pedicels (tight)	Fine	Medium	Even in size
4	Hybrid 7028	Semi-domed	Medium	Short pedicels (tight)	Fine	Medium	Even in size
5	Male 7007	Domed	Small	Short pedicels (tight)	Fine	Large	Even in size
6	Male 7009	Semi-domed	Small	Medium pedicels	Fine	Large	Even in size
7	Male 7022	Very deeply domed	Large	Short pedicels (tight)	Fine	Medium	Even in size
8	Male 7028	Domed	Small	Medium pedicels	Fine	Medium	Even in size
9	Inbred 393- 2-19	Deep domed	Medium	Short pedicels (tight)	Fine	Small	Even in size
10	Inbred 393- 2-47	Semi-domed	Medium	Short pedicels (tight)	Fine	Medium	Even in size
11	Marathon	Deep Domed	Medium	Short pedicels (tight)	Medium	Small	Even in size
12	Pinnacle	Deep Domed	Medium	Medium pedicels	Large	Small	Uneven in size

2.4C. Study #1

Head (At Market Maturity) Anthocyanin Coloration

#	I.D.	Leaf Axils	Leaf Veins	Leaf Blade	Entire Plant	Leaf Petiole
1	Hybrid 7007	Absent	Absent	Absent	Absent	Absent
2	Hybrid 7008	-	-	-	-	-
3	Hybrid 7022	Absent	Absent	Absent	Absent	Absent
4	Hybrid 7028	Absent	Absent	Absent	Absent	Absent
5	Male 7007	Absent	Absent	Absent	Absent	Absent
6	Male 7009	Absent	Absent	Absent	Absent	Absent
7	Male 7022	Absent	Absent	Absent	Absent	Absent
8	Male 7028	Absent	Absent	Absent	Absent	Absent
9	Inbred 393-2-19	Absent	Absent	Absent	Absent	Absent
10	Inbred 393-2-47	Slight Pressure	Absent	Absent	Absent	Absent
11	Marathon	Absent	Absent	Absent	Absent	Absent
12	Pinnacle	Absent	Absent	Absent	Absent	Absent

**2.4D. Study #1**  
**Head (At Market Maturity)**

#	I.D.	Color of Head Leaves	Secondary Heads	Prominence of Secondary Heads	Number of Secondary Heads
1	Hybrid 7007	Green	Completely Absent	Weak	0
2	Hybrid 7008	-	-	-	-
3	Hybrid 7022	Green	Completely Absent	Weak	0
4	Hybrid 7028	Green	Basal	Weak	3
5	Male 7007	Green	Completely Absent	Weak	0
6	Male 7009	Green	Basal	Weak	4
7	Male 7022	Green	Completely Absent	Weak	0
8	Male 7028	Green	Completely Absent	Weak	0
9	Inbred 393-2-19	Green	Completely Absent	Weak	0
10	Inbred 393-2-47	Green	Completely Absent	Weak	0
11	Marathon	Green	Auxiliary along entire main stem up to main head	Weak	3
12	Pinnacle	Green	Basal	Weak	1

2.5. Study #1  
Flower Color

#	I.D.	Flower Color	Flower Stalk Color
1	Hybrid 7007	Yellow	Green
2	Hybrid 7008	-	-
3	Hybrid 7022	Yellow	Green
4	Hybrid 7028	Yellow	-
5	Male 7007	Yellow	-
6	Male 7009	Yellow	-
7	Male 7022	Yellow	Green
8	Male 7028	Yellow	-
9	Inbred 393- 2-19	Yellow	Green
10	Inbred 393- 2-47	Yellow	Green
11	Marathon	Yellow	Green
12	Pinnacle	Yellow	Green

2.6. Study #1  
Resistance\*

#	I.D.	Downey Mildew	Buttoning	Blindness	Bolting	Brown Beads	Drought	Cold	Hollow stem	Riceyness	Whiptail
1	Hybrid 7007	9	8	8	5	9	7	5	8	9	9
2	Hybrid 7008	-	-	-	-	-	-	-	-	-	-
3	Hybrid 7022	9	8	8	4	8	7	3	8	9	9
4	Hybrid 7028	9	8	8	4	9	7	4	8	9	9
5	Male 7007	9	8	8	3	9	8	3	9	9	9
6	Male 7009	9	8	8	5	9	8	3	9	9	9
7	Male 7022	-	8	8	6	8	8	3	8	9	9
8	Male 7028	9	8	8	5	9	8	3	8	9	9
9	Inbred 393- 2-19	9	8	8	6	9	8	6	9	9	9
10	Inbred 393- 2-47	9	8	8	5	9	8	5	9	9	9
11	Marathon	3	8	8	7	8	7	7	8	9	9
12	Pinnacle	3	8	8	6	5	5	4	8	5	9

\* 1 = Most susceptible  
5 = Intermediate  
9 = Most resistant



**2.7. Study #1**  
**Heat Tolerance\***

#	I.D.	Heat Tolerance*
1	Hybrid 7007	9
2	Hybrid 7008	-
3	Hybrid 7022	8
4	Hybrid 7028	9
5	Male 7007	9
6	Male 7009	8
7	Male 7022	7
8	Male 7028	8
9	Inbred 393- 2-19	8
10	Inbred 393- 2-47	8
11	Marathon	2
12	Pinnacle	4

- \* 1 = Most susceptible  
5 = Intermediate  
9 = Most tolerant

### Comparative Analysis Study #2

In a second study (Study #2) various broccoli lines were analyzed and characterized for heat tolerance. Daily high and low temperature measurements for study #2 are presented in Table 3.

5 As in study #2, the daily temperatures were generally quite warm and on some days hot.

In study #2, the following broccoli lines were analyzed: Hybrid 7007, Hybrid 7009, Hybrid 7028, Male 7007, Male 7009, Male 7028, Inbred 393-2-19, Inbred 393-2-47, Marathon, Pinnacle,  
10 98-2061, 98-2088, Inbred 393-2-32 and 4267-1. 98-2061 results from a cross between 393-2-19 and 1551 (393-2-19/1551). As indicated above, 1551 is the same derivative as 7881. The line 98-2088 results from across between 393-2-19 and M7009 (393-2-19/1551). Line 2192 is derived from the same line as 4267-1.

15 In study #2, broccoli seeds were sown in the greenhouse on April 27. Broccoli seedlings were transferred to the field on June 13.

The comparative data collected in study #2 are shown in Table 4.

20 Table 4.1 shows the length of the harvest period, the plant and head height at harvest, the type of plant branches and the plant habit at harvest. Of particular relevance is that the broccoli plants of this invention have a significantly longer harvest period than the commercially available hybrids Marathon and Pinnacle. A longer  
25 harvest period offers growers greater flexibility in harvesting and therefore greatly reduces costs.

Tables 4.2A-4.2C show characteristics of outer leaves at harvest. Tables 4.3A-4.3B and 4.4A show characteristics of the harvested broccoli heads. Table 4.5 shows heat tolerance data.

30 Of particular relevance is the data in Table 4.5, which shows

that the broccoli plants of this invention are heat tolerant whereas the commercially available varieties are not.

[illegible]

**TABLE 3**

**Temperatur Data for Study #2**

Date	Temperature (°F)		
	Max	min	Average
08/08	75	57	63
08/09	81	60	66
08/10	77	58	65
08/11	78	57	65
08/12	79	53	63
08/13	78	55	63
08/14	83	54	64
08/15	77	56	62
08/16	73	56	63
08/17	89	53	63
08/18	83	54	66
08/19	82	59	69
08/20	77	59	66
08/21	87	59	69
08/22	85	56	70
08/23	83	59	69
08/24	82	62	70
08/25	83	57	68
08/26	83	57	68
08/27	83	51	70
08/28	83	56	69
08/29	84	59	69
08/30	82	55	67
08/31	83	59	70
09/01	84	59	70
09/02	81	56	68
09/03	85	59	69
09/04	95	59	73
09/05	87	58	70
09/06	80	55	65
09/07	88	53	66
09/08	86	59	69
09/09	82	55	66
09/10	80	54	66
09/11	79	58	67
09/12	78	54	65
09/13	78	53	63
09/14	79	54	65
09/15	80	56	66

2020-09-15 10:00:00

**TABLE 3**

**Temperature Data for Study #2**

Date	Temperature (°F)		
	Max	min	Average
09/16	85	51	68
09/17	79	54	66
09/18	78	54	64
09/19	82	48	63
09/20	88	51	68
09/21	89	51	66
09/22	87	49	66
09/23	102	54	75
09/24	97	59	74
09/25	87	61	72
09/26	80	57	67
09/27	87	52	68
09/28	95	52	73
09/29	90	59	69
09/30	94	54	65
10/01	74	56	63
10/02	76	57	64
10/03	83	51	66
10/04	81	51	65
10/05	83	50	64
10/06	69	49	59
10/07	70	46	57
10/08	72	43	57
10/09	65	52	60
10/10	62	44	53
10/11	68	44	55
10/12	74	44	57
10/13	83	42	61
10/14	89	44	65
10/15	95	49	68
10/16	96	50	68
10/17	87	50	64
10/18	81	46	59
10/19	64	49	56
10/20	70	49	57
10/21	74	45	55
10/22	70	45	54
10/23	68	49	57
10/24	72	42	55
10/25	75	38	56

2006-09-16 09:00

### Temperature Data for Study #2

Date	Temperature (°F)		
	Max	min	Average
10/26	79	40	57
10/27	75	42	56

**TABLE 4 COMPARATIVE ANALYSIS Study #2**

**4.1. Maturity: Main Crop at 50% Harvest/Plant At Harvest**

#	I.D.	Length of Harvest Period (Days)	Plant Height (inches)	Head Height (inches)	Plant Branches	Plant Habit
1	Hybrid 7007	5	30	22	Medium	Spreading
2	Hybrid 7009	5	28 $\frac{1}{4}$	22	Medium	Spreading
3	Hybrid 7028	4	25 $\frac{5}{8}$	17	Medium	Spreading
4	Male 7007	3	36	33	Medium	Spreading
5	Male 7009	4	37	25 $\frac{1}{2}$	Many	Spreading
6	Male 7028	-	31	23 $\frac{1}{2}$	Medium	Spreading
7	Inbred 393-2-19	6	27	20	Medium	Spreading
8	Inbred 393-2-47	4.5	26	20	Few	Spreading
9	Marathon	1	30.5	20	Many	Intermediate
10	Pinnacle	1	30	26 $\frac{3}{4}$	Medium	Intermediate
11	98-2061	5	26	18	Medium	Intermediate
12	98-2088	4	29 $\frac{1}{4}$	23 $\frac{3}{4}$	Medium	Intermediate
13	Inbred 393-2-32	6	27	18 $\frac{1}{4}$	Medium	Intermediate
14	98-2192	6	28	16	Medium	Spreading

4.2A. Study #2  
Outer Leaves (At Harvest)

#	I.D.	# Leaves Per Plant	Leaf Width (inches)	Leaf Length (inches)	Petiole Length (inches)	Length/Width Ratio
1	Hybrid 7007	23	8	18 1/4	5	2:1
2	Hybrid 7009	26	8	17	5	2:1
3	Hybrid 7028	22	6 1/2	16	4 3/4	2:1
4	Male 7007	24	10	19	7	2:1
5	Male 7009	32	9	21 1/2	6 3/4	2:1
6	Male 7028	18	10 1/2	22 1/2	6 1/2	2:1
7	Inbred 393-2-19	21	7	16 1/2	5 1/2	2:1
8	Inbred 393-2-47	17	5 3/4	11 3/4	3 1/2	2:1
9	Marathon	32	7	18	8 1/4	2:1
10	Pinnacle	25	5 5/8	14 1/2	7 1/4	2:1
11	98-2061	19	6 1/2	16 1/2	6 1/4	2:1
12	98-2088	28	7 1/4	14 3/4	3 1/2	2:1
13	Inbred 393-2-32	21	6 1/2	19	8 1/4	2:1
14	98-2192	28	8 1/4	18 1/2	7 3/4	2:1



4.2B. Study #2  
Outer Leaves (At Harvest)

#	I.D.	Leaf Attachment	Wax Presence	Foliage Color	Leaf Shape	Leaf Base
1	Hybrid 7007	Petiolate	Strong	Medium green	Elliptic	Pointed
2	Hybrid 7009	Petiolate	Strong	Medium green	Elliptic	Blunt
3	Hybrid 7028	Petiolate	Strong	Medium green	Elliptic	Blunt
4	Male 7007	Petiolate	Strong	Medium green	Elliptic	Blunt
5	Male 7009	Petiolate	Strong	Medium green	Broad elliptic	Blunt/pointed
6	Male 7028	Petiolate	Strong	Medium green	Broad elliptic	Blunt/pointed
7	Inbred 393-2-19	Petiolate	Strong	Medium green	Elliptic	Blunt
8	Inbred 393-2-47	Petiolate	Strong	Medium green	Elliptic	Blunt
9	Marathon	Petiolate	Strong	Blue-green	Elliptic	Blunt
10	Pinnacle	Petiolate	Strong	Medium green	Narrow elliptic	Blunt
11	98-2061	Petiolate	Strong	Dark green	Elliptic	Blunt
12	98-2088	Petiolate	Strong	Medium green	Narrow elliptic	Blunt
13	Inbred 393-2-32	Petiolate	Strong	Medium green	Elliptic	Blunt
14	98-2192	Petiolate	Strong	Dark green	Elliptic	Blunt

4.2C. Study #2  
Outer Leaves (At Harvest)

#	I.D.	Leaf Apex	Leaf Margins	Leaf Veins	Attitude	Torsion	Profile
1	Hybrid 7007	Blunt	Slightly wavy	Intermediate	Horizontal	Weak	Planar
2	Hybrid 7009	Blunt	Slightly wavy	Intermediate	Semi-erect/erect	Weak	Planar
3	Hybrid 7028	Blunt	Slightly wavy	Thin	Semi-erect/erect	Weak	Planar
4	Male 7007	Blunt	Slightly wavy	Intermediate	Semi-erect/erect	Weak	Planar
5	Male 7009	Blunt	Very wavy	Intermediate	Semi-erect/erect	Weak	Planar/convex
6	Male 7028	Blunt	Slightly wavy	Thick	Horizontal/semi-erect	Weak	Planar
7	Inbred 393-2-19	Blunt	Slightly wavy	Intermediate	Semi-erect	Weak	Concave
8	Inbred 393-2-47	Blunt	Slightly wavy	Intermediate	Horizontal/semi-erect	Weak	Concave/planar
9	Marathon	Blunt	Very wavy	Intermediate	Horizontal	Intermediate	Concave
10	Pinnacle	Blunt	Slightly wavy	Intermediate	Semi-erect/erect	Intermediate	Convex
11	98-2061	Blunt	Slightly wavy	Thin	Semi-erect	Weak	Planar
12	98-2088	Blunt	Slightly wavy	Intermediate	Semi-erect/erect	Weak	Concave
13	Inbred 393-2-32	Blunt	Slightly wavy	Intermediate	Horizontal	Weak	Planar
14	98-2192	Blunt	Slightly wavy	Intermediate	Semi-erect/erect	Weak	Planar/convex

4.3A. Study #2.  
Head (At Market Maturity)

#	I.D.	Head Diameter (inches)	Head Depth (inches)	Head Weight (gm)	Color	Head Shape
1	Hybrid 7007	10	5 1/4	904.9	Blue/green	Transverse elliptic
2	Hybrid 7009	7	3 1/4	306.4	Purple/ blue/green	Transverse elliptic
3	Hybrid 7028	4 3/4	3	117	Blue/green	Transverse elliptic
4	Male 7007	4 1/4	2 1/2	85.6	Medium green	Circular
5	Male 7009	3 3/4	2 1/4	103.6	Dark green/ Blue/green	Transverse broad elliptic
6	Male 7028	6	3 1/2	450.3	Light purple/ dark green	Transverse elliptic
7	Inbred 393-2-19	5	3	176.3	Blue/green purple	Transverse elliptic narrow
8	Inbred 393-2-47	4 3/4	2 3/4	136.4	Light green/ purple	Transverse elliptic
9	Marathon	4 1/2	2 1/4	313.1	Yellow	Transverse elliptic
10	Pinnacle	6 1/4	4 3/4	336.3	Blue/green	Transverse elliptic
11	98-2061	5 1/2	3	184.1	Blue/green	Transverse broad elliptic
12	98-2088	5 3/4	3 1/2	184.1	Blue/green/ purple	Transverse elliptic
13	Inbred 393-2-32	3 1/4	2 1/2	67.2	Medium green/ blue/green	Transverse broad elliptic
14	98-2192	5 1/2	3	226.0	Blue/green	Transverse elliptic



4.4A. Study #2  
Head (At Market Maturity)

#	I.D.	Color of Head Leaves	Secondary Heads	Prominence of Secondary Heads	# of Secondary Heads
1	Hybrid 7007	---	---	---	---
2	Hybrid 7009	---	---	---	---
3	Hybrid 7028	---	Basal	Weak	1
4	Male 7007	---	Basal	Weak	0
5	Male 7009	Green	---	---	---
6	Male 7028	---	---	---	---
7	Inbred 393-2-19	---	---	---	---
8	Inbred 393-2-47	---	---	---	---
9	Marathon	---	Completely absent	Weak	0
10	Pinnacle	---	Basal	Weak	3
11	98-2061		Basal	Weak	1
12	98-2088		Basal	Intermediate	4
13	Inbred 393-2-32		---	---	---
14	98-2192		Combination	Intermediate	3

**4.5. Study #2**  
**Heat Tolerance**

#	I.D.	Heat Tolerance*
1	Hybrid 7007	7
2	Hybrid 7009	8
3	Hybrid 7028	7
4	Male 7007	---
5	Male 7009	7/8
6	Male 7028	---
7	Inbred 393-2-19	5/6
8	Inbred 393-2-47	5/6
9	Marathon	2
10	Pinnacle	1
11	98-2061	7/8
12	98-2088	7
13	Inbred 393-2-32	---
14	98-2192	8/9

\*      1 = Most susceptible  
         5 = Intermediate  
         9 = Most tolerant

**Comparative Analysis Study #3**

In a third study (Study #3) various broccoli lines were analyzed and characterized for heat tolerance. Lines tested included H7009, H7007, H7028, H7010 H7021R, Marathon, Pinnacle, etc.

5 Lines which include a backslash (/) between the two lines represent a cross between the two lines. The second line on the right side of the backslash, is the "male" line in the cross. The "male" line in the cross is the source of the pollen in the cross. For example 393-2-19/7770 represents a cross between 393-2-19 and 7770 wherein  
10 7770 was the source of the pollen and 393-2-19 was the recipient of the pollen. Single plant selections were made of the crosses. The resulting seed was then selfed. The data presented is summary data based upon an entire row of plants.

As indicated above, H7021R results from a cross between  
15 5580 and 393-2-19 (5580/393-2-19)

Daily high and low temperature measurements for study #3 are presented in Table 5. As in studies #'s 1 and 2, the growth temperatures during study #3 were generally quite warm and sometimes hot.

20 Various broccoli lines were analyzed for heat tolerance. The heat tolerance data is presented in Table 6.

The commercial hybrids (Marathon, Pinnacle, Premium Crop, Patriot, Laguna, Monte Cristo, Greenbelt, Everest, CMS Liberty, and Landmark) averaged a score of 2.83 for heat tolerance. The new  
25 heat tolerant hybrids (7007, 7009, and 7028) that are the subject of this patent application averaged 7.00 for heat tolerance. As discussed above, the heat tolerance scale goes from one (1) to nine (9), with one (1) the most susceptible and nine (9) very highly resistant as described above. In general, ratings of five (5) or below  
30 are unmarketable in a heat stress growth condition and represent

significant economic loss to the broccoli growers if such a level of heat stress reaction occurs in their broccoli fields.

The broccoli lines from study #3 were also comparatively analyzed for bead size, yield, head shape, extension and maturity.

5 The results are presented in Table 7.

10 The bead size rating is on a scale of 1-5. A bead size rating of 5.0 represents very, very large beads. A bead size rating of 1.0 represents very, very small beads such as cauliflower beads. A bead size rating of 4.0 represents large beads. An ideal bead size rating is 3.0 to 4.0 with a maximum desirable rating of 3.7 to 3.8. A combination of bead size rating of 3.8 to 4.0 combined with a high bead size uniformity rating is also acceptable.

15 The yield rating is on a scale of 1 to 10 where a rating of 10 represents a maximum estimated yield for a particular trial. A yield rating of 8.0 compared to a yield rating of 7.0 represents an approximate increase in yield of 50%. While high yields are generally desirable, at the highest yield ratings, hollow core may develop undesirably. A combination of high yield, high heat tolerance (and, therefore head-holding ability) good extension and good uniformity are most desirable. The heat tolerant lines and hybrids of the invention generally exhibited high yields. The high yielding capacity of these lines and hybrids is thought to be due, in part, to an ability to keep increasing head size while maintaining desirable commercial characteristics under heat stress.

25 The head shape, extension and maturity ratings were on a 0 to 10 scale. Head shape is an important selection criterion for broccoli. Head shape ratings of 7.0 to 8.0 are most desired. A head shape score of 3.0 represents a completely flat to nearly concave head. A head shape score of 4.0 to 5.0 represents a small head not yet approaching a semi-dome. A head shape score of 6.0 represents a semi-dome shaped head. A head shape score of 7.0 represents a

30



good, solid dome. A head shape score of 8.0 represents a deep dome. A head shape score of 9.0 represents a very pointed dome in the shape of a Christmas tree.

5 Head extension is also an important selection criterion. Head extension is a comparative measurement of the distance between the broccoli head itself and the leaves surrounding the broccoli. If the head is surrounded by leaves, the head is difficult to harvest. Ideally, the head will be extended up above the leaves to permit easy harvesting of the broccoli. An extension rating of 3.0 represents a head that is buried fairly deep within the leaf canopy. A head extension rating of 5.0 represents a plant having a head which extends only slightly above the leaf canopy. A head extension value of 7.0 represents significant extension of the broccoli head out of the canopy. Commercially available broccoli line Marathon has a head extension rating of 6.5-7.0. The best tolerant broccoli lines of this invention have an extension rating of generally around 7.3. An extension rating of less than 5.0 is undesirable because the head is surrounded by too many leaves making the broccoli difficult to harvest.

20 Maturity is also an important selection criterion. The smaller the maturity rating number the earlier the harvest date. The larger the maturity rating number the later the harvest date. A late harvest date is indicative of a line which takes longer to reach maturity and, therefore, longer to produce a commercially acceptable head.

25 Generally, a smaller maturity rating number is preferred because the broccoli grower is able to harvest his/her crop sooner. Later maturing lines (with higher maturity rating numbers) are acceptable so long as they continue to produce commercially acceptable heads with a proper head size, coloring, head shape, etc.

Prior art commercially available line Marathon generally has a maturity rating of 5.5/6.0. In contrast, the heat tolerant broccoli lines of this invention generally have a maturity rating of around 6.2.

00660000 062204  
T02250 20069800

**TABLE 5**

**Temperature Data for Study #3**

Date	Temperature (°F)		
	Max	Min	Average
07/03	79	47	62
07/04	84	48	63
07/05	75	52	60
07/06	75	52	62
07/07	75	52	61
07/08	71	48	59
07/09	63	54	58
07/10	70	54	59
07/11	73	53	59
07/12	74	52	59
07/13	85	53	65
07/14	87	54	69
07/15	82	50	63
07/16	72	32	62
07/17	76	56	64
07/18	83	58	68
07/19	89	52	69
07/20	83	53	67
07/21	88	53	71
07/22	100	55	78
07/23	99	59	77
07/24	88	56	69
07/25	95	54	72
07/26	81	58	70
07/27	76	55	63
07/28	78	55	62
07/29	75	56	62
07/30	72	56	61
07/31	72	57	62
08/01	82	57	65
08/02	83	56	65
08/03	88	54	68
08/04	83	56	66
08/05	77	56	64
08/06	74	58	64
08/07	79	59	66
08/08	90	56	71

**TABLE 5**

**Temperature Data for Study #3**

Date	Temperature (°F)		
	Max	Min	Average
08/09	98	59	74
08/10	109	60	81
08/11	100	61	78
08/12	91	58	70
08/13	83	56	66
08/14	84	52	64
08/15	81	50	62
08/16	86	50	66
08/17	92	53	71
08/18	98	58	75
08/19	97	60	75
08/20	92	57	71
08/21	89	58	68
08/22	74	54	61
08/23	74	53	61
08/24	72	51	61
08/25	69	53	60
08/26	71	52	60
08/27	77	54	63
08/28	81	50	63
08/29	86	51	67
08/30	87	50	67
08/31	83	51	64
09/01	69	54	59
09/02	81	51	61
09/03	91	48	64
09/04	95	52	71
09/05	94	46	72
09/06	95	57	71
09/07	91	55	69
09/08	93	54	71
09/09	98	54	69
09/10	91	58	71
09/11	83	61	69
09/12	89	64	73
09/13	95	68	77
09/14	93	64	75
09/15	77	56	64

**TABLE 5**

**Temperature Data for Study #3**

Date	Temperature (°F)		
	Max	Min	Average
09/16	68	56	60
09/17	83	32	70
09/18	85	52	67
09/19	87	55	69
09/20	88	57	68
09/21	76	55	62
09/22	73	53	61
09/23	72	52	60
09/24	71	51	59
09/25	65	54	59
09/26	65	49	57
09/27	71	46	46
09/28	73	52	60
09/29	64	48	57
09/30	70	56	59
10/01	63	56	58
10/02	69	47	59
10/03	69	44	55
10/04	76	43	58
10/05	83	44	63
10/06	87	47	65
10/07	78	45	60
10/08	74	51	61
10/09	72	43	57
10/10	75	45	57
10/11	78	40	56
10/12	73	46	58
10/13	75	45	59
10/14	69	32	59
10/15	71	40	54
10/16	74	43	57
10/17	77	39	57

09/16 09/17 09/18 09/19 09/20 09/21 09/22 09/23 09/24 09/25 09/26 09/27 09/28 09/29 09/30 10/01 10/02 10/03 10/04 10/05 10/06 10/07 10/08 10/09 10/10 10/11 10/12 10/13 10/14 10/15 10/16 10/17

**TABLE 6**

**Comparative Analysis: Study #3**

#	ID	Heat Tolerance Rating (0-9)**
1	Marathon	1
2	Pinnacle	3
4	H7009	8
5	Premium Crop	4-
6	H7007	7+
8	H7028	6
9	Patriot	1
10	H7010	
11	Laguna	5
12	H7021R	5-
13	Montecristo	4.1
14	Greenbelt	3-
15	393-2-19/7770	5-
16	393-2-19/7778	6-
17	393-2-19/7861	6
18	Sultan	2.5
19	Tierra	6-
20	393-2-19/7864	
21	393-2-19/7865	4+
22	Everest	2
23	Liberty	2+
24	Marathon	3
25	7881/M7007	7-
26	7770-2/393-2-47	6
27	7770/7935	6
28	7770/7935	6
29	7770/7887	7-
30	Landmark	2
31	H7009	7
32	8092/7825	3
33	8092/7795	4+
34	8092/7883	
35	8030/7935	
36	8030/7914	

7770-2-19/7861

**TABLE 6**

**Comparative Analysis: Study #3**

#	ID	Heat Tolerance Rating (0-9)**
37	H7007	7-
38	Pinnacle	2 +
39	Greenbelt	2
40	393-2-19/1692	7-
41	393-2-19/1524	

\*\* 0 = Most susceptible  
5 = Intermediate  
9 = Most tolerant

2025-06-24 10:23:50

### Comparative Analysis: Study #3

5



#### Comparative Analysis Study #4

In a fourth study, various broccoli lines were analyzed and characterized for heat tolerance. Daily high and low temperature measurements for study #3 are presented in Table 8. The maximum daily temperatures during study #4 were generally cooler than the maximum daily temperatures of study #3. In the fourth study, the maximum daily temperature was never greater than 92°F. In contrast, during study #3, the maximum daily temperature was greater than 95°F on several days.

The heat tolerance ratings for several broccoli lines analyzed during study #4 are present in Table 9. The heat tolerant broccoli lines of the invention consistently exhibited heat tolerance ratings of around 7.0. In contrast, the heat tolerant ratings for the commercially available lines for study #4 averaged around 5.0. The heat tolerance ratings for the commercially available lines were generally higher in study #4 than study #3 because the temperatures were cooler during study #4 than in study #3. Since the commercially available lines were exposed to generally cooler temperatures in study #4, the heat tolerance ratings for these lines were higher during study #4.

In study #4, the heat tolerant broccoli lines were also compared to commercially available lines regarding yield, bead size, head shape, extension and uniformity. The rating scale is the same as that for comparative study #3, Table 7. Uniformity represents a comparative measurement of the similarity between the various plants within a line. High uniformity is desired by growers because it allows them to maximize their harvest efficiency. A minimum uniformity rating of 6.0 to 6.5 is generally viewed as commercially acceptable. A rating of 8.0 represents highly uniform broccoli lines. Uniformity values less than 6.0 are generally viewed as commercially unacceptable. The results are presented in Table 10.

In addition to being heat tolerant, the lines of this invention consistently gave higher head shape and extension ratings than the commercially available lines.

2025-01-20 10:00:00

### Temperature Data for Study #4

Date	Temperature (°F)	
	Max	Min
05/01	66	50
05/02	65	49
05/03	66	52
05/04	70	41
05/05	83	43
05/06	79	49
05/07	76	51
05/08	75	42
05/09	74	41
05/10	79	42
05/11	74	50
05/12	75	54
05/13	79	42
05/14	68	46
05/15	71	42
05/16	76	42
05/17	82	46
05/18	78	48
05/19	69	51
05/20	67	52
05/21	82	46
05/22	81	49
05/23	75	52
05/24	80	52
05/25	80	53
05/26	75	51
05/27	78	51
05/28	75	51
05/29	61	52
05/30	73	53
05/31	75	51
06/01	70	52
06/02	65	53
06/03	68	48
06/04	68	51
06/05	80	52
06/06	80	53

**TABLE 8**

**Temperature Data for Study #4**

Date	Temperature (°F)	
	Max	Min
06/07	75	43
06/08	77	44
06/09	75	44
06/10	81	48
06/11	81	47
06/12	85	51
06/13	84	52
06/14	82	58
06/15	80	57
06/16	84	52
06/17	84	49
06/18	88	51
06/19	84	52
06/20	84	50
06/21	79	54
06/22	86	53
06/23	83	52
06/24	85	54
06/25	83	52
06/26	83	49
06/27	95	49
06/28	99	54
06/29	98	55
06/30	87	51
07/01	87	52
07/02	85	52
07/03	87	52
07/04	87	52
07/05	87	52
07/06	87	52
07/07	87	51
07/08	88	52
07/09	88	52
07/10	90	52
07/11	90	52
07/12	90	53
07/13	91	55
07/14	90	55

06/06/2006 10:30:00

**TABLE 8**

**Temperature Data for Study #4**

Date	Temperature (°F)	
	Max	Min
07/15	87	53
07/16	86	54
07/17	88	54
07/18	89	54
07/19	89	54
07/20	89	54
07/21	89	53
07/22	89	53
07/23	87	54
07/24	87	54
07/25	89	54
07/26	89	54
07/27	89	53
07/28	89	54
07/29	90	54
07/30	88	54
07/31	90	54
08/01	90	54
08/02	88	54
08/03	88	54
08/04	88	54
08/05	90	53
08/06	91	54
08/07	90	55
08/08	90	54
08/09	90	55
08/10	88	55
08/11	85	55
08/12	86	54
08/13	85	54
08/14	86	54
08/15	87	54
08/16	88	54
08/17	87	54
08/18	86	54
08/19	86	54
08/20	86	53
08/21	85	52

### Temperature Data for Study #4

Date	Temperature (°F)	
	Max	Min
08/22	85	53
08/23	88	53
08/24	87	53
08/25	87	53
08/26	87	54
08/27	88	53
08/28	89	53
08/29	89	53
08/30	86	54
08/31	87	54
09/01	89	53
09/02	88	53
09/03	88	53
09/04	88	54
09/05	88	54
09/06	86	54
09/07	86	54
09/08	86	53
09/09	86	53
09/10	86	53
09/11	86	53
09/12	85	52
09/13	84	53
09/14	85	53
09/15	85	52
09/16	84	52
09/17	84	52
09/18	83	52
09/19	84	51
09/20	84	51
09/21	86	50
09/22	86	50
09/23	85	51
09/24	86	52
09/25	84	53
09/26	82	51
09/27	83	53
09/28	83	51

[illegible]

**TABLE 8**

**Temperature Data for Study #4**

Date	Temperature (°F)	
	Max	Min
09/29	84	52
09/30	85	51
10/01	84	50
10/02	82	50
10/03	83	50
10/04	83	50
10/05	84	50
10/06	82	50
10/07	80	49
10/08	80	48
10/09	81	49
10/10	80	48
10/11	79	48
10/12	79	48
10/13	80	49
10/14	79	48
10/15	78	47
10/16	78	47
10/17	80	46
10/18	78	46
10/19	76	47
10/20	75	48
10/21	76	47
10/22	77	46
10/23	75	47
10/24	77	46
10/25	75	48
10/26	74	46
10/27	74	45
10/28	71	45
10/29	73	44
10/30	72	43
10/31	73	43

### Heat Tolerance Data for Study #4

#	ID	Heat Tolerance Rating*
1	4243-1	7.1 -
2	4263-1	7.2
3	4267-1	7.0
4		7.2
5	4274-1	6.9
6	4274-2	6.9
7	4278-1	7.4
8	4284-1	7.2
9	4285-1	6.8
10	4308-2	6.8
11	4309-1	6.7
12	4318-1	6.3
13	4320-1	7.0
14	4320-2	7.0
15	4321-1	7.1
16	4354-1	7.2
17	4354-2	7.2
18	4355-1	6.5
19	4377-1	7.1
20	4395-2	6.5
21	4412-1	6.9
22	4430-1	7.4
23	4432-1	6.9
24	4437-1	7.0
25	4450-1	6.6
26	4450-2	6.6
27	4460-1	7.1
28	4462-1	7.2
29	4465-1	7.3
30	4476-1	7.1
31	Pinnacle	5.0
32	Marathon	5.0
33	Greenbelt	4.5
34	7007	7.0
35	4201	7.0
36	4208	3.0
37	4209	6.0
38	4212	6.1
39	4219	7.0
40	4221	6.9



**TABLE 9**

**Heat Tolerance Data for Study #4**

#	ID	Heat Tolerance Rating*
41	4237	6.2
42	4280	6.1
43	4287	6.1
44	4288	7.3
45	4289	6.0
46	4290	6.7
47	4291/4459	7.2
48	4301	6.5
49	4303	7.1
50	4304	7.1
51	4317	6.4
52	4338	4.9
53	4370	6.2
54	4415	6.7
55	4418	5.5
56	4441	6.3
57	4442	6.3
58	4468	6.8
60	4470	6.5

- \*      0 = Most susceptible  
         5 = Intermediate  
         9 = Most tolerant

20060309 1620

### Yield, Bead Size, Head Shape, Extension and Uniformity Data for Study #4

#	ID	Yield	Bead Size	Head Shape	Extension	Uniformity
1	4243-1	8.8	3.6	7.0	5.5	6.5
2	4263-1	7.0	3.4	7.3	7.3	6.8
3	4267-1	6.8	3.9	6.8	7.1	6.9
4	4267-1	6.9	3.9	7.1	7.2	
5	4274-1	7.2	3.5	7.3	7.5	6.8
6	4274-2	7.2	3.5	7.3	7.5	6.8
7	4278-1	7.6	3.7	7.4	7.2	6.8
8	4284-1	7.3	3.5	7.4	7.5	6.9
9	4285-1	7.4	3.7	7.3	7.3	6.0
10	4308-2	7.0	3.8	6.7	7.9	6.5
11	4309-1	6.9	3.7	6.5	8.2	6.4
12	4318-1		3.6	7.5	7.6	6.9
13	4320-1	7.0	3.6	7.5	7.5	7.0
14	4320-2	7.0	3.6	7.5	7.5	7.0
15	4321-1	7.0	3.7	7.6	7.2	6.9
16	4354-1	7.2	3.6	7.4	7.2	7.1
17	4354-2	7.2	3.6	7.4	7.2	7.1
18	4355-1	6.8	3.9	6.1	7.8	6.5
19	4377-1	7.0	3.7	7.4	7.0	6.8
20	4395-2	7.0	3.7	6.8	7.2	6.8
21	4412-1	7.0	3.8	6.5	7.8	6.1
22	4430-1	7.1	3.3	7.0	7.6	6.9
23	4432-1	7.4	3.5	7.2	7.5	7.0
24	4437-1	7.3	3.6	7.4	7.3	6.1

**TABLE 10**

**Yield, Bead Size, Head Shape, Extension  
and Uniformity Data for Study #4**

#	ID	Yield	Bead Size	Head Shape	Extension	Uniformity
25	4450-1	7.5	3.6	7.1	7.2	6.5
26	4450-2	7.5	3.6	7.1	7.2	6.5
27	4460-1	7.3	3.7	7.1	7.0	7.0
28	4462-1	7.2	3.5	7.3	7.3	6.7
29	4465-1	7.4	3.5	7.2	7.1	6.2
30	4476-1	7.8	3.5	7.4	7.1	6.5
31	Pinnacle	7.2	3.6	6.0	7.2	6.9
32	Marathon	7.3	3.3	6.0	7.0	6.7
33	Greenbelt	6.9	3.5	5.0	7.0	6.9
34	7007	7.5	3.8	7.3	7.1	7.0

**TRANSGENIC BROCCOLI**

The broccoli varieties of this invention can be transformed with useful genes to make heat tolerant transgenic broccoli varieties.

- 5 Such useful genes include "terminator genes", herbicide resistant genes, insect resistant genes, virus resistant genes and the like.

To introduce isolated genes or a group of genes into the genome of plant cells such as broccoli an efficient host gene vector system is necessary. The foreign genes should be expressed in the transformed plant cells and consistently transmitted (somatically and sexually) to the next generation of cells produced. The vector should be capable of introducing, maintaining and expressing a gene in plant cells, from a variety of sources, including but not limited to plants and animals, bacteria, fungi, yeast or virus. Additionally it should be possible to introduce the vector into a wide variety of plants. The location of the new gene in the plant genome may be

10

15

## Directed genetic modification and expression of foreign genes

Using recombinant DNA techniques and bacterial genetics, foreign pieces of DNA can be inserted into T-DNA in *Agrobacterium*.

Transformation of broccoli is well known in the art of molecular biology. For example, in Cao, et al. "Transgenic broccoli with high levels of *Bacillus thuringiensis* CryIC protein control diamondback moth larvae resistant to CryIA or CryIC," *Molecular Breeding* 5:131-141 (1999), discloses one method of generation of transgenic broccoli. Such procedures can readily and easily be followed to produce transgenic heat tolerant broccoli plants.

### ISOLATION OF HEAT TOLERANCE GENES

Now that heat tolerant broccoli plants have been identified and isolated, the identification of the gene or genes involved in heat tolerance is a straightforward process. One of ordinary skill in the art can identify genes involved in heat tolerance by comparing the DNA of heat tolerant and heat sensitive broccoli plants. One such method of isolating heat tolerance gene is the use of a matrix mill available from Cornell University in Ithaca, New York. The use of such a device greatly facilitates the isolation of heat tolerant genes.

The device is capable of breaking up 96 small tissue samples simultaneously in sodium hydroxide, releasing the tissue's DNA and denaturing the protein. After the extraction, the tissue sample is then neutralized and the DNA is simultaneously diluted. Once diluted, the DNA is ready for analysis. Using the matrix mill one can compare several heat tolerant to several heat sensitive broccoli lines simultaneously.

In addition to using the matrix mill, basic molecular biological techniques may be utilized by one of ordinary skill in the art to isolate the heat tolerant broccoli gene. Such procedures are outlined in detail in Ausubel, et al. (Eds) (1987) "Current Protocols in Molecular Biology," John Wiley and Sons, New York.

Once the heat tolerant gene or genes are identified in broccoli, the corresponding heat tolerant gene or genes can be isolated in other plants through various hybridization techniques as described in Ausubel, et al.

Furthermore, biological material can be isolated from the seeds and plants of this invention by procedures well known in the art. Such material may include but is not limited to DNA, RNA, protein and carbohydrates.

**DEPOSIT INFORMATION**

Representative of, but not limiting the invention, Applicants have deposited seeds from M7028, M7007, M7009 and 393-2-19 with the American Type Culture Collection.

- 5 Applicants have made available to the public without restriction a deposit of at least 2500 seeds of broccoli M7028 with the American Type Culture Collection (ATCC), Rockville, MD 20852 which has been assigned ATCC number 203530.

- 10 Applicants have made available to the public without restriction a deposit of at least 2500 seeds of broccoli M7007 with the American Type Culture Collection (ATCC), Rockville, MD 20852 which has been assigned ATCC number 203531.

- 15 Applicants have made available to the public without restriction a deposit of at least 2500 seeds of broccoli M7009 with the American Type Culture Collection (ATCC), Rockville, MD 20852 which has been assigned ATCC number 203532.

- 20 Applicants have made available to the public without restriction a deposit of at least 2500 seeds of broccoli 393-2-19 with the American Type Culture Collection (ATCC), Rockville, MD 20852 which has been assigned ATCC number 203533.

- 25 The deposits will be maintained in the ATCC depository, which is a public depository, for a period of 30 years, or 5 years after the most recent request, or for the effective life of the patent, whichever is longer, and will be replaced if a deposit becomes nonviable during that period.

- 30 Although the foregoing invention has been described in some detail by way of illustration and examples for purposes of clarity and understanding, it will be obvious that certain modifications and alternative embodiments of the invention are contemplated which do not depart from the spirit and scope of the invention as defined by the foregoing teachings and appended claims.